

Parametric Duality and Kernelization: Lower Bounds and Upper Bounds on Kernel Size*

JIANER CHEN[†] HENNING FERNAU[‡] IYAD A. KANJ[§] GE XIA[¶]

Abstract

Determining whether a parameterized problem is kernelizable and has a small kernel size has recently become one of the most interesting topics of research in the area of parameterized complexity and algorithms. Theoretically, it has been proved that a parameterized problem is kernelizable if and only if it is fixed-parameter tractable. Practically, applying a data-reduction algorithm to reduce an instance of a parameterized problem to an equivalent smaller instance (i.e, a kernel) has led to very efficient algorithms, and now goes hand-in-hand with the design of practical algorithms for solving \mathcal{NP} -hard problems. Well-known examples of such parameterized problems include the VERTEX COVER problem, which is kernelizable to a kernel of size bounded by $2k$, and the PLANAR DOMINATING SET problem, which is kernelizable to a kernel of size bounded by $335k$. In this paper we develop new techniques to derive upper and lower bounds on the kernel size for certain parameterized problems. In terms of our lower bound results, we show, for example, that unless $\mathcal{P} = \mathcal{NP}$, PLANAR VERTEX COVER does not have a problem kernel of size smaller than $4k/3$, and PLANAR INDEPENDENT SET and PLANAR DOMINATING SET do not have kernels of size smaller than $2k$. In terms of our upper bound results, we further reduce the upper bound on the kernel size for the PLANAR DOMINATING SET problem to $67k$, improving significantly the $335k$ previous upper bound given by Alber et al. This latter result is obtained by introducing a new set of reduction and coloring rules, which allows the derivation of nice combinatorial properties in the kernelized graph leading to a tighter bound on the size of the kernel. The paper also shows how this improved upper bound yields a simple and competitive algorithm for the PLANAR DOMINATING SET problem.

Keywords. parameterized algorithm, planar graph, dominating set, vertex cover, independent set, kernel

1 Introduction

Many practical algorithms for \mathcal{NP} -hard problems start by applying data reduction subroutines to the input instances of the problem. The hope is that after the data reduction phase the instance of the problem has shrunk to a moderate size. This makes the applicability of a second phase, such as a branch-and-bound phase, to the resulting instance more feasible. Weihe showed in [41]

*An extended abstract of this paper appeared in STACS 2005.

[†]Department of Computer Science, Texas A&M University, College Station, TX 77843, USA. This research was supported in part by the NSF under grants CCF-0430683 and CCR-0311590. Email: chen@cs.tamu.edu.

[‡]FB 4-Abteilung Informatik, Universität Trier, D-54286 Trier, Germany.
Email:fernau@informatik.uni-trier.de.

[§]The corresponding author. School of CTI, DePaul University, 243 S. Wabash Avenue, Chicago, IL 60604, USA. Email: ikanj@cs.depaul.edu. This work was supported in part by DePaul University Competitive Research Grant.

[¶]Department of Computer Science, Acopian Engineering Center, Lafayette College, Easton PA 18042, USA. Email: gexia@cs.lafayette.edu.

how a practical pre-processing algorithm for a variation of the DOMINATING SET problem, called the RED/BLUE DOMINATING SET problem, resulted in breaking up input instances of the problem into much smaller instances. Abu-Khzam et al. [2], in their implementation of algorithms for the VERTEX COVER problem, observed the following: “In many cases, reduction was so effective that it eliminated the core completely, and with it the need for decomposition and search”. Similar success was reported with DOMINATING SET as well in [3].

On the other hand, many applications seek solutions of very small sizes to fairly large input instances of \mathcal{NP} -hard problems. This has been the main concern for the area of parameterized computation. A *parameterized problem* is a set of instances of the form (x, k) , where x is the input instance, and k is a nonnegative integer called the *parameter*. A parameterized problem is said to be *fixed-parameter tractable* [17] if there is an algorithm that solves the problem in time $f(k)|x|^c$, where c is a fixed constant and $f(k)$ is a recursive function. The development of efficient parameterized algorithms has provided a new approach for practically solving problems that are theoretically intractable. For example, parameterized algorithms for the \mathcal{NP} -hard problem VERTEX COVER [9, 13] have found applications in biochemistry [10] and variants thereof are applicable to problems arising in chip manufacturing [11, 21, 24], and parameterized algorithms in computational logic [35] have provided an effective method for solving practical instances of the ML TYPE-CHECKING problem, which is complete for the class EXPTIME [30].

The notion of a parameterized problem being parameterized tractable, and of the problem having a good data reduction algorithm, turn out to be very closely related. Informally speaking, a *kernelization*—precisely defined below, is a data reduction procedure that reduces an instance of the problem to another (smaller) instance called the *kernel*. It has been proved that a parameterized problem is fixed-parameter tractable if and only if the problem is kernelizable [18].

Designing efficient parameterized algorithms and constructing kernels of reasonable sizes have been two of the main topics of research in the area of parameterized computation recently. More specifically, constructing a problem kernel has become one of the main components in the design of an efficient parameterized algorithm for a problem [9, 11, 12, 13], and designing efficient parameterized algorithms for a parameterized problem now goes hand-in-hand with the construction of a problem kernel of a moderate size for the problem. Two of the most celebrated problems that have been receiving a lot of attention recently from both perspectives, are the VERTEX COVER and PLANAR DOMINATING SET problems. After a long sequence of algorithms, the VERTEX COVER problem can be solved in time $O(1.274^k + kn)$ [14]. Moreover, the VERTEX COVER problem enjoys a kernel of size bounded by $2k$, and reducing this bound further seems to be a very challenging task, since it would probably lead to an approximation algorithm for the problem of ratio smaller than 2—a result believed by many people to be unlikely. The PLANAR DOMINATING SET problem as well has undergone some extensive study which culminated in a recent algorithm solving the problem in time $O(2^{15.13\sqrt{k}} + n^3)$ [25]. Recently, and after many strenuous efforts, it was shown that the PLANAR DOMINATING SET problem has a problem kernel of size $335k$ that is computable in $O(n^3)$ time [5]. The question of whether such a bound on the problem kernel could be significantly improved remained open.

In this paper we develop new techniques to derive upper and lower bounds on the kernel size for certain parameterized problems. We define the notion of *duality* of a parameterized problem. Many parameterized tractable problems are the dual of parameterized intractable problems (see [19, 34, 38]). As an example, consider the VERTEX COVER and INDEPENDENT SET problems. If n denotes the number of vertices in the whole graph G , then it is well-known that (G, k) is a YES-instance of VERTEX COVER if and only if (G, k_d) , where $k_d = n - k$, is a YES-instance of INDEPENDENT SET. In this sense, INDEPENDENT SET is the parametric dual problem of VERTEX

COVER. While VERTEX COVER is fixed-parameter tractable on general graphs, INDEPENDENT SET is not [17]. Similarly, while DOMINATING SET is fixed-parameter intractable on general graphs, its parametric dual, called NONBLOCKER, is fixed-parameter tractable, see [15]. The landscape changes when we turn our attention towards special graph classes, e.g., problems on *planar graphs* [6]. Here, for example, both INDEPENDENT SET and DOMINATING SET are fixed-parameter tractable. In fact, and in contrast to what was stated above, there are quite many problems for which both the problem itself and its dual are fixed-parameter tractable. This is also true for problems on graphs of bounded genus, as well as on graphs of bounded degree.

The beauty of problems which are together with their dual problems fixed-parameter tractable is that this constellation allows from an algorithmic standpoint for a two-sided attack on the original problem. This two-sided attack enabled us to derive lower bounds on the kernel size for such problems (under classical complexity assumptions). For example, we show that unless $\mathcal{P} = \mathcal{NP}$, PLANAR VERTEX COVER does not have a kernel of size smaller than $4k/3$, and PLANAR INDEPENDENT SET and PLANAR DOMINATING SET do not have kernels of size smaller than $2k$. To the authors' knowledge, this is the first group of results establishing lower bounds on the kernel size of parameterized problems. We also show that some lower bound results obtained using the techniques devised in this paper are sharp by exhibiting a family of graph classes on which the lower bound on the kernel size of the restricted \mathcal{NP} -hard VERTEX COVER problem approaches the upper bound $2k$ with an arbitrary precision.

Whereas the lower bounds on the kernel size for PLANAR VERTEX COVER and PLANAR INDEPENDENT SET come close to the known upper bounds of $2k$ and $4k$ on the kernel size for the two problems, respectively, the lower bound derived for PLANAR DOMINATING SET is still very far from the $335k$ upper bound on the problem kernel, which was given by Alber et al. [5]. To bridge this gap, we investigate the problem of finding a kernel of smaller size for the PLANAR DOMINATING SET problem, and derive better upper bounds on the problem kernel for the problem. We improve the reduction rules proposed in [5], and introduce new rules that *color* the vertices of the graph enabling us to observe many new combinatorial properties of its vertices. These properties allow us to prove a much stronger bound on the number of vertices in the reduced graph. We show that the PLANAR DOMINATING SET problem has a kernel of size $67k$ that is computable in $O(n^3)$ time. This is a significant improvement over the results in [5]. We finally show how the resulting bound on the kernel size yields a very simple algorithm for the PLANAR DOMINATING SET problem that beats some previous algorithms for the problem, and whose running time comes even close to some of the recently proposed algorithms.

2 Preliminaries

A graph G is said to be *planar* if G can be embedded on the plane such that no two edges in G cross. It is well-known that deciding whether a graph is planar, and constructing a planar embedding of the graph in such case, can be done in linear time [31]. The number of edges in a planar graph with n vertices for $n \geq 3$ is bounded by $3n - 6$ [16].

A dominating set in a graph G is a set of vertices D such that every vertex in G is either in D or adjacent to at least one vertex in D . The size of a dominating set D is the number of vertices in D . A *minimum dominating set* of G is a dominating set with the minimum size. We will denote by $\gamma(G)$ the size of a minimum dominating set in G . The PLANAR DOMINATING SET problem, abbreviated PLANAR-DS henceforth, is the following: given a planar graph G and a positive integer k , either construct a dominating set for G of size at most k , or report that no such a dominating set exists.

It is well-known that the PLANAR-DS problem is \mathcal{NP} -complete [27].

A *parameterized problem* P is a subset of $\Sigma^* \times \mathbb{N}$, where Σ is a fixed alphabet and \mathbb{N} is the set of all non-negative integers. Therefore, each instance of the parameterized problem P is a pair (I, k) , where the second component k is called the *parameter*. The language $L(P)$ is the set of all YES-instances of P . We say that the parameterized problem P is *fixed-parameter tractable* [17] if there is an algorithm that decides whether an input (I, k) is a member of $L(P)$ in time $f(k)|I|^c$, where c is a fixed constant and $f(k)$ is a recursive function independent of the input length $|I|$. The class of all fixed parameter tractable problems is denoted by FPT.

A mapping $s : \Sigma^* \times \mathbb{N} \rightarrow \mathbb{N}$ is called a *size function* for a parameterized problem P if:

- $0 \leq k \leq s(I, k)$,
- $s(I, k) \leq |I|$, and
- $s(I, k) = s(I, k')$ for all appropriate k, k' (*independence*). Hence, we can also write $s(I)$ for $s(I, k)$.

A problem P together with its size function s are denoted (P, s) . The *dual problem* P_d of P is the problem whose corresponding language (i.e., the set of YES-instances) $L(P_d) = \{(I, s(I) - k) \mid (I, k) \in L(P)\}$. The dual of the dual of a problem (with a given size function) is again the original problem. We give some examples below.

d -HITTING SET

Given: A hypergraph $G = (V, E)$ with *edge degree* bounded by d , i.e., $\forall e \in E, |e| \leq d$

Parameter: k

Question: Is there a *hitting set* of size at most k , i.e.,

$$\exists C \subseteq V, |C| \leq k, \forall e \in E, C \cap e \neq \emptyset?$$

The special case in which $d = 2$ corresponds to the VERTEX COVER problem in undirected graphs. Let $L(d\text{-HS})$ denote the language of d -HITTING SET. Taking as size function $s(G) = |V|$, it is clear that the dual problem obeys $(G, k_d) \in L(d\text{-HS}_d)$ if and only if G has an *independent set* of cardinality k_d .

DOMINATING SET

Given: A (simple) graph $G = (V, E)$

Parameter: k

Question: Is there a *dominating set* of size at most k , i.e.,

$$\exists D \subseteq V, |D| \leq k, \forall v \in V \setminus D \exists w \in D, (w, v) \in E?$$

Taking as size function $s(G) = |V|$, it is clear that the dual problem obeys $(G, k_d) \in L(\text{DS}_d)$ if and only if G has a *nonblocker set* (i.e., the complement of a dominating set) of cardinality k_d .

Generally speaking, it is easy to “correctly” define the dual of a problem for the so-called selection problems as formalized in [7]. The concept of duality is less clear say for weighted graph problems (with the slight exception of ROMAN DOMINATION; see [22]). Also, different graph parameterizations like treewidth seem to possess no natural dualization.

A *kernelization (reduction)* for a parameterized problem P with size function s is a polynomial-time computable reduction which maps an instance (I, k) onto (I', k') such that: (1) $s(I') \leq g(k)$ (g is a recursive function), (2) $k' \leq k$, and (3) $(I, k) \in L(P)$ if and only if $(I', k') \in L(P)$. I' is called the *problem kernel* of I . It is known (see [18]) that a parameterized problem is fixed-parameter tractable if and only if it has a kernelization. Of special interest to us in this paper are problems with *linear kernels* in which $g(k) = \alpha k$ for some constant $\alpha > 0$. Such small kernels are known for many important graph problems restricted to planar graphs.

3 Lower bounds on kernel size

Practice in the study of parameterized algorithms has suggested that improved kernelization can lead to improved parameterized algorithms. Many efforts have been made towards obtaining smaller kernels for well-known \mathcal{NP} -hard parameterized problems (see for example [5, 13, 18]). A natural question to ask along this line of research is about the limit of polynomial time kernelization. In this section we develop techniques for deriving lower bounds on the kernel size for certain well-known \mathcal{NP} -hard parameterized problems.

3.1 General lower bound results

Theorem 3.1 *Let (P, s) be an \mathcal{NP} -hard parameterized problem (with size function s). Suppose that P admits an αk kernelization and its dual P_d admits an $\alpha_d k_d$ kernelization, where $\alpha, \alpha_d \geq 1$. If $(\alpha - 1)(\alpha_d - 1) < 1$ then $\mathcal{P} = \mathcal{NP}$.*

PROOF. Suppose that the assumption of the theorem is true, and let $r(\cdot)$ denote the assumed linear kernelization reduction for P . Similarly, let $r_d(\cdot)$ be the linear kernelization reduction for P_d . Consider the following reduction R , which on input (I, k) of P performs the following:

if $k \leq \frac{\alpha_d}{\alpha + \alpha_d} s(I)$ **then** compute $r(I, k)$;
else compute $r_d(I, s(I) - k)$.

Let (I', k') be the instance computed by the reduction R . If $k \leq \frac{\alpha_d}{\alpha + \alpha_d} s(I)$, then $s(I') \leq \alpha k \leq \frac{\alpha \alpha_d}{\alpha + \alpha_d} s(I)$. Otherwise:

$$\begin{aligned} s(I') &\leq \alpha_d k_d \\ &= \alpha_d (s(I) - k) \\ &< \alpha_d \left(s(I) - \frac{\alpha_d}{\alpha + \alpha_d} s(I) \right) \\ &= \frac{\alpha \alpha_d}{\alpha + \alpha_d} s(I). \end{aligned}$$

Since $(\alpha - 1)(\alpha_d - 1) < 1$, or equivalently $\frac{\alpha \alpha_d}{\alpha + \alpha_d} < 1$, by repeatedly applying R (at most polynomially-many times), the problem P can be solved in polynomial time. This completes the proof. \square

The condition “ $\alpha, \alpha_d \geq 1$ ” in the previous theorem is not crucial in the light of the following lemma.

Lemma 3.2 *Let (P, s) be a parameterized problem such that P admits a kernelization $r(\cdot)$ with $s(r(I, k)) \leq \alpha k$ for some $\alpha < 1$. Then P is in \mathcal{P} .*

PROOF. According to our definition of the size function, we have $s(I') \geq k'$ for each instance (I', k') . This is in particular true for the parameter k' of the problem kernel instance $I' = r(I, k)$. Therefore, $k' \leq \alpha k$ for some $\alpha < 1$. By repeated kernelization, we arrive at a problem with arbitrarily small parameter, and hence of arbitrarily small size. In fact, we need $\mathcal{O}(\log k)$ many such kernelizations, each of them requiring polynomial time. It follows that the given problem can be decided in polynomial time. \square

REMARK. The assumption $s(I) \geq k$ is crucial here. As a concrete “counterexample,” consider the *decision tree problem*, specified by n objects $X = \{x_1, \dots, x_n\}$, t boolean tests $T = \{T_1, \dots, T_t\}$ and a parameter k . In this setting, a decision tree is a binary tree B whose leaves are (uniquely) labeled with objects and whose inner nodes are labeled with tests such that on the path from the root to the leaf labeled x_i , tests are performed that uniquely distinguish x_i from all other objects. The overall length of all paths from the root to each leaf is usually considered as the cost function. The question is if there exists a decision tree with cost bounded by k . This problem is known to be \mathcal{NP} -complete (see [32]).

If $n = 2^\ell$, the decision tree with optimal cost is surely the complete binary tree (possibly not attainable with the given set of tests), since it is optimally balanced. Hence, we have $k > n \log_2 n$ (otherwise, an algorithm can simply answer NO); this can be seen as a trivial kernelization algorithm. Therefore, $n \in o(k)$. This can be interpreted as giving the (to our knowledge) first natural parameterized problem having a sub-linear kernel. On the other hand, this relation also implies that $s(I, k) < k$ is true here, so that the previous lemma does not lead to a contradiction with the known \mathcal{NP} -hardness result.

The problem here is the seemingly innocuous choice of the size function as being $n = |X|$. Observe that any “reasonable” encoding of an instance would rather use $n \log n$ bits, since each element of X would need to get a name. This way, the problem would disappear. \square

3.2 Concrete lower bound results

From Theorem 3.1, and assuming $\mathcal{P} \neq \mathcal{NP}$, we immediately obtain the following list of corollaries.

- **Corollary 3.3** *For any $\epsilon > 0$, there is no $(4/3 - \epsilon)k$ kernel for PLANAR VERTEX COVER.*

PROOF. The four-color theorem implies a $4k$ -kernelization for PLANAR INDEPENDENT SET, which is the dual problem of PLANAR VERTEX COVER. \square

- **Corollary 3.4** *For any $\epsilon > 0$, there is no $(2 - \epsilon)k$ kernel for PLANAR INDEPENDENT SET. This result remains true if we restrict the problem to graphs of maximum degree bounded by three, or even to planar graphs of maximum degree bounded by three (both problems are \mathcal{NP} -hard).*

PROOF. The general VERTEX COVER problem, which is the dual of the INDEPENDENT SET problem, has a $2k$ -kernelization [13] (based on a result of Nemhauser and Trotter). This kernelization is both planarity and bounded-degree preserving. \square

- **Corollary 3.5** *For any $\epsilon > 0$, there is no $(3/2 - \epsilon)k$ -kernelization for VERTEX COVER restricted to triangle-free planar graphs (this problem is still \mathcal{NP} -hard [40, Chapter 7]).*

PROOF. Based on a theorem by Grötzsch (which can be turned into a polynomial-time coloring algorithm; see [29]) it is known that planar triangle-free graphs are 3-colorable. This implies a $3k$ kernel for INDEPENDENT SET restricted to this graph class, which gives the result. Observe that the mentioned $2k$ -kernelization for VERTEX COVER on general graphs preserves planarity and triangle-freeness, which implies that this restriction of the problem has a $2k$ -kernelization. \square

- **Corollary 3.6** *For any $\epsilon > 0$, there is no $(335/334 - \epsilon)k_d$ kernel for PLANAR NONBLOCKER.*

PROOF. A $335k$ kernel for PLANAR-DS was derived in [5]. \square

- **Corollary 3.7** *For any $\epsilon > 0$, there is no $(2 - \epsilon)k$ kernel for PLANAR-DS. This remains true when further restricting the graph class to planar graphs of maximum degree three (the problem is still \mathcal{NP} -hard).*

PROOF. In [20], a $2k_d$ -kernelization for NONBLOCKER on general graphs which preserves planarity and degree bounds, was derived (see also [37, Theorem 13.1.3]). \square

Corollary 3.8 (see [15]) *For any $\epsilon > 0$, there is no $(2 - \epsilon)k$ kernel for DOMINATING SET ON CUBIC GRAPHS.*

This is interesting, since that case is the best match between upper and lower bounds for domination problems.

The above results open a new line of research, and prompt us to ask whether we can find examples of problems such that the derived kernel sizes are optimal (unless $\mathcal{P} = \mathcal{NP}$), and whether we can close the gaps between the upper bounds and lower bounds on the kernel size. According to our previous discussion, PLANAR VERTEX COVER on triangle-free graphs is our “best match:” we know how to derive a kernel of size $2k$, and (assuming $\mathcal{P} \neq \mathcal{NP}$) we know that no kernel smaller than $3k/2$ exists. On the other hand, the $335k$ upper bound on the kernel size for PLANAR-DS [5] is very far from the $2k$ lower bound proved above. In the next sections, we improve this upper bound to $67k$ in an effort to bridge the huge gap between the upper bound and lower bound on the kernel size for this problem. This allows us to state:

Corollary 3.9 *Assuming $\mathcal{P} \neq \mathcal{NP}$, there is no $(67/66 - \epsilon)k_d$ kernel for PLANAR NONBLOCKER for any choice of $\epsilon > 0$.*

REMARK. Since “Euler-type” theorems exist for graphs of bounded genus g , it can be shown that there is a constant c_g such that each graph of genus g is c_g -colorable. Therefore, lower bounds on the kernel sizes of VERTEX COVER on graphs of genus g can be derived. For triangle-free graphs of genus g , Thomassen has shown that the corresponding constant c'_g is in $\mathcal{O}(g^{1/3}(\log g)^{-2/3})$ (see [28, 39]). \square

REMARK. Recently, Fomin and Thilikos [26] were able to extend the linear kernel result for DOMINATING SET to graphs on surfaces of bounded genus. Therefore, our lower bound results extend to these more general graph classes as well. \square

3.3 Can we improve on the lower bounds?

In the following, we reproduce a construction that is essentially due to Paul Seymour.¹ This construction shows that the lower bound results obtained using the techniques devised in this section can be sharp for certain problems.

Consider the following family \mathcal{G}_n of graph classes. A graph G is in \mathcal{G}_n if and only if it satisfies the following two conditions.

1. $G = (V, E)$ can be partitioned into $2n + 1$ mutually disjoint independent sets, i.e., $V = I_1 \cup \dots \cup I_{2n+1}$, $I_i \cap I_j = \emptyset$ for all $1 \leq i < j \leq 2n + 1$, and the induced graphs $G[I_i]$ contain no edges.
2. The edge set E can be partitioned into $2n + 1$ groups E_1, \dots, E_{2n+1} such that

$$E_i = E(G[I_i \cup I_{i \bmod (2n+1)+1}]).^2$$




Figure 1: An example of a graph in \mathcal{G}_n .

Figure 1 provides an example of a graph in \mathcal{G}_n .

Since each of these classes is closed under taking induced subgraphs, we can deduce by the Nemhauser-Trotter kernelization [13]:

Lemma 3.10 VERTEX COVER restricted to \mathcal{G}_n admits a kernel of size $2k$ (within \mathcal{G}_n).

Since the graphs in \mathcal{G}_n are “nearly bipartite,” we have the following result.

Lemma 3.11 INDEPENDENT SET restricted to \mathcal{G}_n admits a kernel of size $(2 + 1/n)k_d$ (within \mathcal{G}_n).

PROOF. Let $G = (V, E) \in \mathcal{G}_n$ with an independent set decomposition $V = I_1 \cup \dots \cup I_{2n+1}$ that

¹Personal communication (12.07.05).

²We assume that the graph G is given with a certificate of membership in \mathcal{G}_n , which is a partitioning of its vertex set into the $2n + 1$ subsets.

certifies this membership. To simplify the notation, we assume that additions and subtractions of indices are all performed modulo $2n + 1$. Consider the sets:

$$J_i = I_i \cup I_{i+2} \cup \dots \cup I_{i-3} \quad 1 \leq i \leq 2n + 1.$$

J_i greedily collects “every second” set starting at I_i so that each set J_i forms an independent set of G . It can be easily verified that:

$$\sum_{i=1}^{2n+1} |J_i| = n|V|.$$

This shows that there exists an index i for which the set J_i contains at least a fraction $n/(2n+1)$ of all the vertices. Moreover, such a J_i can be found in polynomial time. Therefore, we can answer YES straightaway whenever we are given a graph $G = (V, E) \in \mathcal{G}_n$ with a parameter $k_d \leq (n/(2n+1))|V|$, as an instance of INDEPENDENT SET. This means that we have $|V| < (2 + 1/n)k_d$ for all the remaining instances. \square

Theorem 3.12 *For each n , VERTEX COVER restricted to \mathcal{G}_n is \mathcal{NP} -complete.*

PROOF. Membership in \mathcal{NP} is inherited from the general VERTEX COVER problem. We will show that 3-SAT is polynomial-time reducible to VERTEX COVER restricted to \mathcal{G}_n . We highlight the main elements in the reduction here and leave the verification of some of the details to the interested reader.

Let $\mathcal{C} = \{C_1, \dots, C_m\}$ be a collection of clauses. Without loss of generality, we can assume that $|C_i| = 3$ for all $1 \leq i \leq m$. Let ℓ_i^j refers to the j th literal in clause C_i , i.e., $\ell_i^j = y_i^j$ or $\ell_i^j = \bar{y}_i^j$ for some variable $y_i^j \in X = \{x_1, \dots, x_r\}$.

We construct a graph $G = (V, E) \in \mathcal{G}_n$ as follows. For each variable x_i , we introduce a cycle $(v_i^1, \dots, v_i^{4n+2})$ of even length. Clearly, $2n + 1$ of these vertices will be in any feasible vertex cover. For each clause C_i , we introduce a cycle $(u_i^1, \dots, u_i^{2n+1})$ of odd length. Clearly, $n + 1$ of these vertices will be in any feasible vertex cover. Summarizing, the graph described so far will have at least $r(2n + 1) + m(n + 1)$ vertices in any feasible cover. Since we will now add more edges to this graph, the lower bound on the size of the vertex cover will still be valid. At the same time, we will maintain the property that $I_j = \{v_i^j, v_i^{j+2n+1} \mid 1 \leq i \leq r\} \cup \{u_i^j \mid 1 \leq i \leq m\}$, where $1 \leq j \leq 2n + 1$, are all independent sets.

If $\ell_i^1 = y_i^1 = x_q$, then we will make u_i^1 adjacent to v_q^2 . If $\ell_i^2 = y_i^2 = x_q$, then we will make u_i^2 adjacent to v_q^{2n+2} . If $\ell_i^3 = y_i^3 = x_q$, then we will make u_i^3 adjacent to v_q^2 .

If $\ell_i^1 = \bar{y}_i^1 = \bar{x}_q$, then we will make u_i^1 adjacent to v_q^{2n+3} . If $\ell_i^2 = \bar{y}_i^2 = \bar{x}_q$, then we will make u_i^2 adjacent to v_q^1 . If $\ell_i^3 = \bar{y}_i^3 = \bar{x}_q$, then we will make u_i^3 adjacent to v_q^{2n+3} .

Now, if x_i is set to true in a satisfying assignment of the given 3-SAT instance, then we put v_i^j into the vertex cover if and only if j is even. If x_i is set to false in a satisfying assignment of the given 3-SAT instance, then we put v_i^j into the vertex cover if and only if j is odd. It is not difficult to verify that a satisfying assignment of \mathcal{C} can be translated into a feasible vertex cover of size $r(2n + 1) + m(n + 1)$.

The same identification of vertices from v_i^j with variable settings allows to translate a feasible vertex cover of size $r(2n + 1) + m(n + 1)$ into a satisfying assignment for the given 3-SAT instance. \square

Corollary 3.13 *Unless $\mathcal{P} = \mathcal{NP}$, the VERTEX COVER problem restricted to \mathcal{G}_n does not have a kernel of size $(2 - \epsilon)k$ for any $\epsilon > 0$.*

PROOF. This follows from Lemma 3.11 and Theorem 3.1. □

REMARK. The above corollary shows that the lower bound results on the kernel size for VERTEX COVER restricted to \mathcal{G}_n obtained using the techniques in this paper are tight. It also shows that it is unlikely that the VERTEX COVER problem on general graphs admits a kernelization of size $(2 - \epsilon)k$ with the property that the produced kernel is a subgraph of the original graph. The reason being that such a kernelization would also be a kernelization for the vertex cover problem restricted to the class \mathcal{G}_n with the same kernel bound. □

3.4 A possible two-sided attack for exact algorithms

With problems having both FPT algorithms for their primal and for their dual parameterizations, we have the possibility of converting both algorithms into a non-parameterized algorithm. This is like attacking the problem from two sides. This means that we can use either of the two FPT algorithms, depending on “to which side” our concrete problem instance is closer.

Theorem 3.14 *Let (P, s) be a parameterized problem and P_d its dual. Assume that both P and P_d are in FPT. Let f be a monotone function. Assume that there is an algorithm A that solves an instance (I, k) of P in time $\mathcal{O}(f(\beta k)p(s(I)))$ for some polynomial p , and that A_d is an algorithm that solves an instance (I, k_d) of P_d in time $\mathcal{O}(f(\beta_d k_d)p_d(s(I)))$ for a polynomial p_d .*

Then, there is an algorithm A' for solving the non-parameterized problem instance I running in time

$$\mathcal{O}\left(f\left(\frac{\beta\beta_d}{\beta + \beta_d}s(I)\right)p'(s(I))\right),$$

for some polynomial p' .

PROOF. The idea is to use algorithm A as long as it is better than using A_d . This means that we have to compare

$$f(\beta k) \quad \text{to} \quad f(\beta_d(s(I) - k_d)).$$

Since f is monotone, this means we can simply compare

$$\beta k \quad \text{to} \quad \beta_d(s(I) - k_d).$$

Some simple algebra shows that we can have the following algorithm A' for the non-parameterized problem P , given an instance I :

for all parameter values k **do**:
if $k \leq \frac{\beta_d}{\beta + \beta_d}s(I)$ **then** compute $A(I, k)$;
else compute $A_d(I, s(I) - k)$;
output the ‘best’ of all computed solutions.

Considering the boundary case $k = \frac{\beta_d}{\beta + \beta_d}s(I)$ gives the claimed worst case running time. Here, $p'(j) = j(p(j) + p_d(j))$. □

Unfortunately, up till now we are lacking good examples that prove this approach superior to published (problem-tailored) exact algorithms.

4 Reduction and coloring rules

In this section we show how to improve the upper bound on the kernel size for the PLANAR-DS problem to $67k$. In the remainder of the paper we will always assume that the graph we are dealing with is planar.

In this section we present an $O(n^3)$ time preprocessing scheme that reduces the graph G to a graph G' , such that $\gamma(G) = \gamma(G')$, and such that given a minimum dominating set for G' , a minimum dominating set for G can be constructed in linear time. We will color the vertices of the graph G with two colors: black and white. Initially, all vertices are colored black. Informally speaking, white vertices will be those vertices that we know for sure when we color them that there exists a minimum dominating set for the graph excluding all of them. The black vertices are all other vertices. Note that it is possible for white vertices to be in some minimum dominating set, but the point is that there exists at least one minimum dominating set that excludes all white vertices. Hence, the black-and-white coloring is only an auxiliary structure and not part of the problem definition. We start with the following definitions that are adopted from [5] with minor additions and modifications.

For a vertex v in G denote by $N(v)$ the set of neighbors of v , and by $N[v]$ the set $N(v) \cup \{v\}$. By removing a vertex v from G , we mean removing v and all the edges incident on v from G . For a vertex v in G , we partition its set of neighbors $N(v)$ into three sets: $N_1(v) = \{u \in N(v) \mid N(u) - N[v] \neq \emptyset\}$; $N_2(v) = \{u \in N(v) - N_1(v) \mid N(u) \cap N_1(v) \neq \emptyset\}$; and $N_3(v) = N(v) - (N_1(v) \cup N_2(v))$. For two vertices v and w we define $N(v, w) = N(v) \cup N(w)$ and $N[v, w] = N[v] \cup N[w]$. We partition $N(v, w)$ into three sets: $N_1(v, w) = \{u \in N(v, w) \mid N(u) - N[v, w] \neq \emptyset\}$; $N_2(v, w) = \{u \in N(v, w) - N_1(v, w) \mid N(u) \cap N_1(v, w) \neq \emptyset\}$; and $N_3(v, w) = N(v, w) - (N_1(v, w) \cup N_2(v, w))$.

Definition 4.1 Let $G = (V, E)$ be a plane graph. A *region* $R(v, w)$ between two vertices v and w is a closed subset of the plane with the following properties:

1. The boundary of $R(v, w)$ is formed by two simple paths P_1 and P_2 in G which connect v and w , and the length of each path is at most three.
2. All vertices that are strictly inside (i.e., not on the boundary) the region $R(v, w)$ are from $N(v, w)$.

For a region $R = R(v, w)$, let $V[R]$ denote the vertices in R , i.e.,

$$V[R] := \{u \in V \mid u \text{ sits inside or on the boundary of } R\}.$$

Let $V(R) = V[R] - \{v, w\}$.

Definition 4.2 A region $R = R(v, w)$ between two vertices v and w is called *simple* if all vertices in $V(R)$ are common neighbors of both v and w , that is, $V(R) \subseteq N(v) \cap N(w)$.

We introduce the following definitions.

Definition 4.3 A region $R = R(v, w)$ between two vertices v and w is called *quasi-simple* if $V[R] = V[R'] \cup R^+$, where $R' = R'(v, w)$ is a simple region between v and w , and R^+ is a set of white vertices satisfying the following conditions:

1. Every vertex of R^+ sits strictly inside R' .
2. Every vertex of R^+ is connected to v and not connected to w , and is also connected to at least one vertex on the boundary of R' other than v .

A vertex in $V(R)$ is called a *simple* vertex, if it is connected to both v and w , otherwise it is called *non-simple*. The set of vertices R^+ , which consists of the non-simple vertices in $V(R)$, will be referred to as $R^+(v, w)$.

For a vertex $u \in V$, denote by $B(u)$ the set of black vertices in $N(u)$, and by $W(u)$ the set of white vertices in $N(u)$. We describe next the reduction and coloring rules to be applied to the graph G . The reduction and coloring rules are applied to the graph until the application of any of them does not change the structure of the graph nor the color of any vertex in the graph. The first two reduction rules, **Rule 1** and **Rule 2**, are slight modifications of Rule 1 and Rule 2 introduced in [5]. The only difference is that in the current paper they are only applied to black vertices, and not to all the vertices as in [5].

Rule 1 ([5]). If $N_3(v) \neq \emptyset$ for some black vertex v , then remove the vertices in $N_2(v) \cup N_3(v)$ from G , and add a new white vertex v' and an edge (v, v') to G .

Rule 2 ([5]). If $N_3(v, w) \neq \emptyset$ for two black vertices v, w , and if $N_3(v, w)$ cannot be dominated by a single vertex in $N_2(v, w) \cup N_3(v, w)$, then we distinguish the following two cases.

Case 1. If $N_3(v, w)$ can be dominated by a single vertex in $\{v, w\}$ then: (1.1) if $N_3(v, w) \subseteq N(v)$ and $N_3(v, w) \subseteq N(w)$, remove $N_3(v, w)$ and $N_2(v, w) \cap N(v) \cap N(w)$ from G and add two new white vertices z, z' and the edges $(v, z), (w, z), (v, z'), (w, z')$ to G ; (1.2) if $N_3(v, w) \subseteq N(v)$ and $N_3(v, w) \not\subseteq N(w)$, remove $N_3(v, w)$ and $N_2(v, w) \cap N(v)$ from G and add a new white vertex v' and the edge (v, v') to G ; and (1.3) if $N_3(v, w) \subseteq N(w)$ and $N_3(v, w) \not\subseteq N(v)$, remove $N_3(v, w)$ and $N_2(v, w) \cap N(w)$ from G and add a new white vertex w' and the edge (w, w') to G .

Case 2. If $N_3(v, w)$ cannot be dominated by a single vertex in $\{v, w\}$, then remove $N_2(v, w) \cup N_3(v, w)$ from G and add two new white vertices v', w' and the edges $(v, v'), (w, w')$ to G .

Rule 3. For each black vertex v in G , if there exists a black vertex $x \in N_2(v) \cup N_3(v)$, color x white, and remove the edges between x and all other white vertices in G .

Rule 4. For every two black vertices v and w , if $N_3(v, w) \neq \emptyset$, then for every black vertex $x \in N_2(v, w) \cup N_3(v, w)$ that does not dominate all vertices in $N_3(v, w)$, color x white and remove all the edges between x and the other white vertices in G .

Rule 5. For every quasi-simple region $R = R(v, w)$ between two vertices v and w , if v is black, then for every black vertex $x \in N_2(v, w) \cup N_3(v, w)$ strictly inside R that does not dominate all vertices in $N_2(v, w) \cup N_3(v, w)$ strictly inside R , color x white and remove all the edges between x and the other white vertices in G .

Rule 6. For every two white vertices u and v , if $N(u) \subseteq N(v)$, and $u \in N_2(w) \cup N_3(w)$ for some black vertex w , then remove v .

Rule 7. For every black vertex v , if every vertex $u \in W(v)$ is connected to all the vertices in $B(v)$, then remove all the vertices in $W(v)$ from G .

Rule 8. For every two black vertices v and w , let $W(v, w) = W(v) \cap W(w)$. If $|W(v, w)| \geq 2$ and

there is a degree-2 vertex $u \in W(v, w)$, then remove all vertices in $W(v, w)$ except u , add a new degree-2 white vertex u' , and connect u' to both v and w .

Figure 2 illustrates **Rules 4, 6, and 8**.

A graph G is said to be *reduced* if every vertex in G is colored white or black, and the application of **Rules 1–8** leaves the graph G unchanged. That is, the application of any of the above rules does not change the color of any vertex in G , nor does it change the structure of G . We have the following theorem.

Theorem 4.1 *Let G be a planar graph with n vertices. Then in time $O(n^3)$ we can construct a planar graph G' from G such that: (1) G' is reduced, (2) $\gamma(G') = \gamma(G)$, (3) there exists a minimum dominating set for G' that excludes all white vertices of G' , and (4) from a minimum dominating set for G' a minimum dominating set for G can be constructed in linear time.*

PROOF. Given a graph G , we first color all its vertices black. We then apply **Rule 1 – Rule 8** given above until the application of any of these rules leaves G unchanged. Let G' be the resulting graph. Then G' is reduced by the definition of a reduced graph. Alber et al [5] noted that each successful application of **Rule 1** and **Rule 2** (i.e., an application that changes the structure of the graph G) reduces the number of vertices in the graph by at least one. Hence, the total number of applications of these two rules is bounded by n . By looking at **Rule 3 – Rule 7**, it is easy to see that each of these rules either reduces the number of vertices in G by at least one, or changes the color of at least one black vertex to white without adding any new vertices to the graph. Moreover, none of **Rule 1 – Rule 7** increases the number of edges in the graph. If we look at **Rule 8**, it is not difficult to see that each successful application of this rule reduces the number of edges in the graph by at least 1. This is true since in a successful application of the rule either $|W(v, w)| > 2$ and in this case the number of vertices and edges decrease after the application of the rule, or $|W(v, w)| = 2$ and there is a vertex in $W(v, w)$ of degree larger than 2 (otherwise the application of the rule does not change the structure of the graph), and hence the removal of $W(v, w)$ decreases the number of edges in the graph. Noting that the number of edges in a planar graph is linear in the number of vertices, and that the application of the rules become unnecessary if the graph does not contain any black vertices, we conclude that the total number of successful applications of the operations in **Rule 1 – Rule 8** is $O(n)$. Alber et al. [5] also showed that **Rule 1** and **Rule 2** can be executed in time $O(n^2)$ when the graph is planar. Similarly, one can show that **Rules 3–8** can also be executed in $O(n^2)$ time (we leave the verification of this simple fact to the interested reader). This, together with the fact that the total number of successful applications of all the rules is $O(n)$, implies that the time needed to construct G' is $O(n^3)$.

To show parts (2) and (3) of the theorem, we prove that after the execution of any of the rules, the resulting graph satisfies conditions (2) and (3) in the theorem. The proof will then follow by an inductive argument on the number of applications of the rules. Denote by H the graph before a rule is executed, and by H' the resulting graph after the rule is executed. Denote by D a minimum dominating set for H excluding all white vertices in H . Initially, $H = G$ and all vertices in H are black. Thus, H trivially satisfies conditions (2) and (3) in the theorem. Suppose now that one of the rules is executed on a graph H satisfying conditions (2) and (3) in the theorem to yield the graph H' . We need to show that H' satisfies conditions (2) and (3) as well.

Suppose that **Rule 1** is executed. The same argument used in [5] shows that $\gamma(H) = \gamma(H')$.³

³The fact that this statement holds true can be easily verified by the reader.




Figure 2: Illustrations of **Rule 4** (top figure), **Rule 6** (middle figure), and **Rule 8** (bottom figure).

What is left is showing that H' has a minimum dominating set consisting only of black vertices. Let D be a minimum dominating set for H consisting of black vertices. Since $N_3(v) \neq \emptyset$, D must contain a vertex in $N_2(v) \cup N_3(v) \cup \{v\}$. If D contains a vertex in $N_2(v) \cup N_3(v)$, then clearly this vertex can be replaced by v which is black. Thus we can assume that D contains v and no vertex in $N_3(v) \cup N_2(v)$. Then D is also a dominating set for H' consisting only of black vertices, and since $\gamma(G) = \gamma(H) = \gamma(H')$, D is a minimum dominating set for H' . It follows that H' satisfies conditions (2) and (3). The proof of **Rule 2** is of the same flavor.

If **Rule 3** is executed, then the black vertices in the set $N_2(v) \cup N_3(v)$, where v is black, will be colored white, and the edges between the white vertices are removed. It suffices to show that after the coloring of one vertex x in $N_2(v) \cup N_3(v)$ white, and removing the edges between x and the other white vertices, conditions (2) and (3) still hold (the same argument can then be applied repetitively to every such vertex). By our inductive hypothesis, before the application of **Rule 3** to H , H had a minimum dominating set D of size equal to $\gamma(G)$ that excludes all white vertices in H . If D contains x , we can substitute x with v and have a minimum dominating set of H consisting only of black vertices in H . Thus, we can assume, without loss of generality, that D does not include x . Since x is the only vertex whose color has changed to white, D consists only of black vertices in H' . Moreover, it is not difficult to see that D is also a dominating set in H' since the edges removed from H are not used to dominate any vertices in H (these edges were incident on vertices that are not in D). Since by removing edges from the graph the size of the minimum dominating set can only increase, we conclude that D is a minimum dominating set for H' excluding all white vertices, and $\gamma(H') = \gamma(H) = \gamma(G)$.

Suppose **Rule 4** is executed. Similarly, we only need to show that conditions (2) and (3) still hold after a vertex x has been colored white. If D contains x , then by the assumption in **Rule 4**, x does not dominate all the vertices in $N_3(v, w)$, and D must also contain at least another vertex x' in $N_2(v, w) \cup N_3(v, w) \cup \{v, w\}$ to dominate $N_3(v, w)$. This is true since only vertices in $N_2(v, w) \cup N_3(v, w) \cup \{v, w\}$ can dominate vertices of $N_3(v, w)$. In such case we can substitute x and x' by v and w and have a minimum dominating set that consists only of black vertices in H . Since x is the only vertex whose color has changed to white, D excludes all white vertices in H' . It is easy to see that the edges that connect white vertices in H are not used by D to dominate any vertex. By an argument similar to the above, it follows that D is a minimum dominating set for H' excluding all white vertices in H' , and $\gamma(H') = |D| = \gamma(H) = \gamma(G)$.

Suppose **Rule 5** is executed and a vertex x is colored white. Let $R = R(v, w)$ be the quasi-simple region that was being processed in the rule, and note that all the vertices in $R^+(v, w)$ are connected to v . Let the boundary of R be (v, y, w, z, v) . Let D be a dominating set for H consisting only of black vertices. If D contains x , then by the assumption in **Rule 5**, x does not dominate all the vertices in $N_2(v, w) \cup N_3(v, w)$ strictly inside R , and D must contain another black vertex $x' \in R(v, w)$ in $N_2(v, w) \cup N_3(v, w) \cup \{v, w, y, z\}$ to dominate the vertices in $N_2(v, w) \cup N_3(v, w)$ that are strictly inside R . Observe that, by the definition of a quasi-simple region, the only vertex that can be dominated by x and not by v is w . We distinguish the following cases:

Case 1. $x' = v$. Since at least one vertex $r \in \{y, w, z\}$ must be black (w is connected to both y and z and no edges exist between white vertices, so it is not possible for all the vertices in $\{y, w, z\}$ to be white) and since all the vertices in $\{y, w, z\}$ dominate w , we can substitute x by r (note that x is dominated by v) to obtain a dominating set consisting of black vertices that excludes x .

Case 2. $x' \neq v$. If x' does not dominate w , then x' must be one of those vertices in R^+ that connect only to v and to the vertices on the boundary of R other than w . Since all such boundary

vertices are dominated by v , and x' is dominated by v as well, we can substitute x' by v in D and the case reduces to **Case 1** above. If x' dominates w , then we can substitute x by v to get a dominating set consisting of black vertices that excludes x .

Thus, we can assume, without loss of generality, that D does not include x . Since x is the only vertex whose color has changed to white, D consists only of black vertices in H' . By an argument similar to above, it follows that D is a minimum dominating set for H' excluding all white vertices in H' , and $\gamma(H') = |D| = \gamma(H) = \gamma(G)$.

Suppose **Rule 6** is executed and a vertex v is removed as described in the rule. Let D be a minimum dominating set for H excluding all white vertices in H . Thus D does not contain v . Since v is the only vertex removed, and no vertices are colored, it follows that D is a dominating set for H' excluding all white vertices in H' . What is left is proving that D is a minimum dominating set for H' . Suppose that H' has a minimum dominating set D' of size strictly smaller than D . Then D' has to cover u , and hence, D' either contains u or a neighbor of u in H' . If D' contains u , since every neighbor of u is also a neighbor of w , and u is a neighbor of w , $(D' \cup \{w\}) - \{u\}$ is a minimum dominating set for H of size smaller than D , a contradiction (note that since w is a neighbor of u , w is a neighbor of v as well, and hence, dominates v). On the other hand, if D' contains a neighbor of u , since $N(u) \subseteq N(v)$, D' also contains a neighbor of v in H , and hence dominates v . Thus, D' is a dominating set for H of size smaller than D , a contradiction. It follows that $|D| = \gamma(H') = \gamma(H) = \gamma(G)$.

Suppose **Rule 7** is executed on a black vertex v , and all vertices in $W(v)$ were removed as described in the rule. Let D be a minimum dominating set for H excluding all white vertices in H . Thus, D does not contain any vertex in $W(v)$. Since the vertices in $W(v)$ are the only vertices that were removed, and no vertices in the graph were colored, it follows that D is a dominating set for H' excluding all white vertices in H' . What is left is proving that D is a minimum dominating set for H' . Suppose that H' has a minimum dominating set D' of size strictly smaller than D . Then D' has to cover v . Hence D' either contains v , or a neighbor of v in $B(v)$ because all the vertices in $W(v)$ were removed. In either case, D' dominates all the removed vertices in $W(v)$ in H , since every vertex in $W(v)$ is adjacent to all vertices in $B(v)$. Therefore D' is also a dominating set for H of size smaller than D , a contradiction. It follows that $|D| = \gamma(H') = \gamma(H) = \gamma(G)$.

To prove the statement for **Rule 8**, let D be a minimum dominating set for H excluding all white vertices in H . Again, D is a dominating set for H' excluding all white vertices in H' . Let D' be a minimum dominating set for H' and suppose, to get a contradiction, that $|D'| < |D|$. Without loss of generality, we can assume that D' contains either v or w (or both), otherwise, to dominate u and u' , D' has to contain both u and u' , which can be substituted by v and w . Now D' is also a dominating set for H of smaller size than D , a contradiction. It follows that D is a minimum dominating set for H' excluding all white vertices in H' , and $\gamma(H') = \gamma(H) = \gamma(G)$.

To prove part (4) of the theorem, note the following: (1) from a minimum dominating set for G' one can construct in $O(n)$ time a minimum dominating set for G' containing only black vertices (this can be achieved by associating, during the reduction phase, with the vertices colored white the black vertices that can replace them); and (2) a minimum dominating set for G' consisting only of black vertices is also a minimum dominating set for G . This completes the proof. \square

5 A problem kernel

Let G be a reduced graph, and let D be a minimum dominating set for G consisting of black vertices such that $|D| = k$. In this section, we will show that the number of vertices n in G is bounded by $67k$. The following definitions are adopted from [5].

Given any dominating set D in a graph G , a D -region decomposition of G is a set \mathfrak{R} of regions between pairs of vertices in D such that:

1. For any region $R = R(v, w)$ in \mathfrak{R} , no vertex in D is in $V(R)$. That is, a vertex in D can only be an endpoint of a region in \mathfrak{R} .
2. No two distinct regions $R_1, R_2 \in \mathfrak{R}$ intersect. However, they may touch each other by having common boundaries.

Note that all the endpoints of the regions in a D -region decomposition are vertices in D . For a D -region decomposition \mathfrak{R} , define $V[\mathfrak{R}] = \bigcup_{R \in \mathfrak{R}} V[R]$. A D -region decomposition is *maximal*, if there is no region R such that $\mathfrak{R}' = \mathfrak{R} \cup R$ is a D -region decomposition with $V[\mathfrak{R}] \subsetneq V[\mathfrak{R}']$.

For a D -region decomposition \mathfrak{R} , associate a planar graph $G_{\mathfrak{R}}(V_{\mathfrak{R}}, E_{\mathfrak{R}})$ with possible multiple edges, where $V_{\mathfrak{R}} = D$, and such that there is an edge between two vertices v and w in $G_{\mathfrak{R}}$ if and only if $R(v, w)$ is a region in \mathfrak{R} . A planar graph with multiple edges is called *thin*, if there is a planar embedding of the graph such that for any two edges e_1 and e_2 between two distinct vertices v and w in the graph, there must exist two more vertices which sit inside the disjoint areas of the plane enclosed by e_1 and e_2 .

Alber et al. [5] showed that the number of edges in a thin graph of n vertices is bounded by $3n - 6$. They also showed that for any plane graph G and a dominating set D of G , there exists a maximal D -region decomposition for G such that $G_{\mathfrak{R}}$ is thin. Since the maximal D -region decomposition in [5] starts with any dominating set D and is not affected by the color a vertex can have, the same results in [5] hold true for our reduced graph G whose vertices are colored black/white, and with a minimum dominating set D consisting only of black vertices. The above discussion is summarized in the following proposition.

Proposition 5.1 *Let G be a reduced graph and D a dominating set of G consisting of black vertices. Then there exists a maximal D -region decomposition \mathfrak{R} of G such that $G_{\mathfrak{R}}$ is thin.*

Corollary 5.2 *Let G be a reduced graph with a minimum dominating set D consisting of k black vertices, and let \mathfrak{R} be a maximal D -region decomposition of G such that $G_{\mathfrak{R}}$ is thin. Then the number of regions in \mathfrak{R} is bounded by $3k - 6$.*

PROOF. The number of regions in \mathfrak{R} is the number of edges in $G_{\mathfrak{R}}$. Since $G_{\mathfrak{R}}$ has $|D| = k$ vertices, by [5], the number of edges in $G_{\mathfrak{R}}$ is bounded by $3k - 6$. \square

In the remainder of this section, \mathfrak{R} will denote a maximal D -region decomposition of G such that $G_{\mathfrak{R}}$ is thin. Let u and v be two vertices in G . We say that u and v are *boundary-adjacent* if (u, v) is an edge on the boundary of some region $R \in \mathfrak{R}$. For a vertex $v \in G$, denote by $N^*(v)$ the set of vertices that are boundary-adjacent to v . Note that for a vertex $v \in D$, since v is black, by **Rule 3**, all vertices in $N_2(v) \cup N_3(v)$ must be white. Note also that, by the definition of a D -region decomposition, all the endpoints of the regions in \mathfrak{R} are vertices in D , and hence are colored black.

Proposition 5.3 *Let $v \in D$. The following are true.*

- (a) (Lemma 6, [5]) Every vertex $u \in N_1(v)$ is in $V[\mathfrak{R}]$.
- (b) The vertex v is an endpoint of a region $R \in \mathfrak{R}$. That is, there exists a region $R = R(x, y) \in \mathfrak{R}$ such that $v = x$ or $v = y$.
- (c) Every vertex $u \in N_2(v)$ which is not in $V[\mathfrak{R}]$ is connected only to v and to vertices in $N^*(v)$.

PROOF. The proof of part (a) appears in [5].

To prove (b), suppose to get a contradiction that v is not the endpoint of any region in \mathfrak{R} . Since $v \in D$, and by the definition of a region, v must be outside every region in \mathfrak{R} . Now v must have a vertex in $N_1(v)$, otherwise, all vertices in $N(v)$ would be white, and hence removed by **Rule 7** (we assume, without loss of generality, that G does not contain any isolated vertices). Let $u \in N_1(v)$. By part (a) above, u must belong to some region $R = R(x, y)$. Observe that u must be on the boundary of R , otherwise v would be a vertex in $V[R]$. Again, by the definition of a region, u is either boundary-adjacent to x or to y . Suppose, without loss of generality, that u is boundary-adjacent to x . But then the degenerated region formed by (x, u, v) does not cross \mathfrak{R} (it only touches R), contradicting the maximality of \mathfrak{R} .

To prove part (c), let u be a vertex in $N_2(v)$ and note that u is white, and suppose that u is connected to a vertex $w \neq v$ such that $w \notin N^*(v)$. Note that w must be in $N_1(v)$ (otherwise w would be white and u and w cannot be adjacent), and hence, by part (a) above, must belong to some region $R = R(x, y)$. Since $u \notin V[\mathfrak{R}]$, w cannot be inside R , and hence, is on the boundary of R . Moreover, by the definition of a region, w must be boundary-adjacent to either x or y . Without loss of generality, assume w is boundary-adjacent to x . Now $w \notin N^*(v)$, so w cannot be boundary-adjacent to v , and $x \neq v$. Consider the degenerated region formed by (v, u, w, x) . This region cannot cross any region in \mathfrak{R} , otherwise it crosses it via (u, w) and u would be in $V[\mathfrak{R}]$. But this contradicts the maximality of \mathfrak{R} since $u \notin V[\mathfrak{R}]$. \square

Let x be a vertex in G such that $x \notin V[\mathfrak{R}]$. Then by part (b) in Proposition 5.3, $x \notin D$. Thus, $x \in N(v)$ for some black vertex $v \in D \subseteq V[\mathfrak{R}]$. By part (a) in Proposition 5.3, $x \notin N_1(v)$, and hence, $x \in N_2(v) \cup N_3(v)$. By **Rule 3**, the color of x must be white. Let $R = R(v, w)$ be a region in $V[\mathfrak{R}]$ of which v is an endpoint (such a region must exist by part (b) of Proposition 5.3). We distinguish two cases.

Case A. $x \in N_3(v)$. Since v is black, by **Rule 1**, this is only possible if $\deg(x) = 1$ and $N_2(v) = \emptyset$ (in this case x will be the white vertex added by the rule). In such case it can be easily seen that we can flip x and place it inside R without affecting the planarity of the graph.

Case B. $x \in N_2(v)$. Note that in this case $N_3(v) = \emptyset$ by **Rule 1** (otherwise $N_2(v) \cup N_3(v)$ would be removed), and x is only connected to v and $N^*(v)$ by part (c) in Proposition 5.3. If $\deg(x) = 2$, by a similar argument to **Case A** above, x can be flipped and placed inside R .

According to the above discussion, it follows that the vertices in G can be classified into two categories: (1) those vertices that are in $V[\mathfrak{R}]$; and (2) those that are not in $V[\mathfrak{R}]$, which are those vertices of degree larger than two that belong to $N_2(v)$ for some vertex $v \in D$, and in this case must be connected only to vertices in $N^*(v)$. To bound the number of vertices in G we need to bound the number of vertices in the two categories. We start with the vertices in category (2).

Let O denote the set of vertices in category (2). Note that all vertices in O are white, and no two vertices u and v in O are such that $N(u) \subseteq N(v)$. To see why the latter statement is true,

note that every vertex in O must be in $N_2(w)$ for some black vertex $w \in D$. So if $N(u) \subseteq N(v)$, then by **Rule 6**, v would have been removed from the graph. To bound the number of vertices in O , we will bound the number of vertices in O that are in $N_2(v)$ where $v \in D$. Let us denote this set by $N^\dagger(v)$. Let $N_\dagger^*(v)$ be the set of vertices in $N^*(v)$ that are neighbors of vertices in $N^\dagger(v)$. Note that every vertex in $N^\dagger(v)$ has degree ≥ 3 , is connected only to v and to $N_\dagger^*(v)$, and no two vertices x and y in $N^\dagger(v)$ are such that $N(x) \subseteq N(y)$.

Assumption 5.4 *For the sake of counting the number of vertices in $N^\dagger(v)$, it is safe to assume that: (1) every vertex in $N^\dagger(v)$ has degree exactly 3; (2) no two vertices x and y in $N^\dagger(v)$ are such that $N(x) \subseteq N(y)$; and (3) vertices in $N^\dagger(v)$ are only connected to v and to vertices in $N_\dagger^*(v)$.*

PROOF. Since properties (2) and (3) are already satisfied by the vertices in $N^\dagger(v)$, we only need to show how we can make the vertices in $N^\dagger(v)$ satisfy property (1) without reducing their number, and without affecting properties (2) and (3). To satisfy property (1), we will remove some edges between vertices in $N^\dagger(v)$ and $N_\dagger^*(v)$ without affecting the other properties. This can be done as follows. List the vertices in $N^\dagger(v)$ in an arbitrary order $\langle u_1, \dots, u_r \rangle$. Start by picking u_1 , then choose any two neighbors of u_1 in $N_\dagger^*(v)$ and remove all edges that join u_1 to all its neighbors other than v and these two chosen neighbors. Inductively, suppose we have processed vertex u_{i-1} , we process vertex u_i as follows. Pick two neighbors w_1^i and w_2^i of u_i in $N_\dagger^*(v)$ such that no vertex in $\{u_1, \dots, u_{i-1}\}$ has both w_1^i and w_2^i as its picked neighbors. Delete all the edges that join u_i to all vertices other than v , w_1^i , and w_2^i . We need to show that it is always possible to carry out this step. Suppose not, and let i be the smallest index such that this is not possible. It is easy to verify using the facts that every vertex in $N^\dagger(v)$ has degree larger than two, and no two vertices x and y are such that $N(x) \subseteq N(y)$, that $i > 3$. Note that it must be the case that $\deg(u_i) > 3$, otherwise, since this step cannot be carried out, the only two neighbors of u_i other than v must be neighbors of some other vertex $u_j \in \{u_1, \dots, u_{i-1}\}$, and hence we would have $N(u_i) \subseteq N(u_j)$ for some $u_j \in \{u_1, \dots, u_{i-1}\}$, contradicting the properties satisfied by category-(2) vertices as discussed above. Let a, b, c be three neighbors of u_i other than v . Since this step cannot be carried out successfully, there must exist three distinct vertices $u_p, u_q, u_s \in \{u_1, \dots, u_{i-1}\}$ such that $\{a, b\} \subset N(u_p)$, $\{a, c\} \subset N(u_q)$, and $\{b, c\} \subset N(u_s)$. Consider the subgraph H of G induced by the set of vertices $\{v, u_p, u_q, u_s, u_i, a, b, c\}$. Then the following is true about the vertices in H : (1) $u_i, u_p, u_q, u_s, a, b, c$ are neighbors of v in H ; (2) v, a, b, c are neighbors of u_i in H ; (3) v, a, b are neighbors of u_p in H ; (4) v, a, c are neighbors of u_q in H ; (5) v, b, c are neighbors of u_s in H ; (6) v, u_i, u_p, u_q are neighbors of a in H ; (7) v, u_i, u_p, u_s are neighbors of b in H ; and (8) v, u_i, u_q, u_s are neighbors of c in H . Using all this information, it is not difficult to verify that H is non-planar (identify vertex a with vertex b along the path (a, u_p, b) to obtain a copy of $K_{3,3}$), contradicting the planarity of G . This completes the proof. \square

Proposition 5.5 $|N^\dagger(v)| \leq 3/2|N_\dagger^*(v)|$.

PROOF. To simplify the counting, by Assumption 5.4, we can assume that: every vertex in $N^\dagger(v)$ has degree exactly 3; no two vertices x and y in $N^\dagger(v)$ are such that $N(x) \subseteq N(y)$; and vertices in $N^\dagger(v)$ are only connected to v and $N_\dagger^*(v)$. Let x be the number of vertices in $N_\dagger^*(v)$, and let $f(x) = |N^\dagger(v)|$. We will show that $f(x) \leq 3/2(x - 1)$. We proceed by induction on x . If $x = 1$, it is clear that $f(x) = 0 \leq 3/2(x - 1)$ since by Assumption 5.4, each vertex in $N^\dagger(v)$ has degree exactly 3. If $x = 2$, then clearly $f(x) \leq 1 \leq 3/2(x - 1)$ since at most one vertex can be connected to v and the two vertices in $N_\dagger^*(v)$ without violating properties (1)–(3) in Assumption 5.4 above.

If $x = 3$, then $f(x) \leq 3$ since at most three vertices can be connected to $N_{\dagger}^*(v)$ without violating properties (1)–(3) in Assumption 5.4 above, each connected to v and to two other vertices in $N_{\dagger}^*(v)$. Inductively, suppose that if $N_{\dagger}^*(v)$ contains y vertices with $3 \leq y < x$, then the number of vertices $f(y)$ in $N^{\dagger}(v)$ satisfies $f(y) \leq 3/2(y - 1)$. Suppose now that $|N_{\dagger}^*(v)| = x$. Let u be a vertex in $N^{\dagger}(v)$, and let a, b be its neighbors in $N_{\dagger}^*(v)$. The vertex u is called *hollow* if the interior of the region enclosed by (u, a, v, b, u) contains no vertices of $N_{\dagger}^*(v)$. If every vertex in $N^{\dagger}(v)$ is hollow, then it is clear that $f(x) \leq x \leq 3/2(x - 1)$ for $x > 3$, and the bound $f(x) = x$ is attained when there are x vertices in $N^{\dagger}(v)$, and every vertex u in $N^{\dagger}(v)$ is adjacent to v and the two neighbors a and b in $N_{\dagger}^*(v)$ immediately to the left and right in the clockwise (or anticlockwise) ordering, respectively, of u in the embedding. Suppose now that there is a vertex $u \in N^{\dagger}(v)$ such that u is not hollow. The edges (u, a) , (u, v) , (u, b) , (v, a) , (v, b) separate the plane into three faces: F_1 enclosed by the cycle (u, a, v, u) , F_2 enclosed by the cycle (u, v, b, u) , and F_3 is the outer face determined by the cycle (u, a, v, b, u) . Let x_1 be the number of vertices in $N_{\dagger}^*(v)$ that are in F_1 including the boundary, x_2 that in F_2 , and x_3 that in F_3 . Note that $1 \leq x_1 < x$ since $a \in F_1$ and $b \notin F_1$, $1 \leq x_2 < x$ since $b \in F_2$ and $a \notin F_2$, and $2 \leq x_3 < x$ since a and b are in F_3 and at least one vertex in $N_{\dagger}^*(v)$ is not in F_3 since u is hollow, and hence, the interior of the face (u, a, v, b, u) contains at least one vertex in $N_{\dagger}^*(v)$. Moreover, $x_1 + x_2 + x_3 = x + 2$, since a and b are the only vertices counted twice when we add the vertices in $N_{\dagger}^*(v)$ that are in F_1 , F_2 , and F_3 . Now every vertex in $N^{\dagger}(v)$ is either: (1) connected to two vertices in $N_{\dagger}^*(v)$ in F_1 , (2) connected to two vertices in $N_{\dagger}^*(v)$ in F_2 , or (3) connected to two vertices in $N_{\dagger}^*(v)$ in F_3 . Note that vertex u satisfies property (3). Since $x_1, x_2, x_3 < x$, by the inductive hypothesis, the number of vertices satisfying (1) is bounded by $f(x_1) \leq 3/2(x_1 - 1)$, the number of vertices satisfying (2) is bounded by $f(x_2) \leq 3/2(x_2 - 1)$, and the number of vertices satisfying (3) is bounded by $f(x_3) \leq 3/2(x_3 - 1)$. Now $f(x) \leq f(x_1) + f(x_2) + f(x_3) \leq 3/2(x_1 + x_2 + x_3) - 9/2 = 3x/2 - 3/2 = 3/2(x - 1)$. This completes the proof. \square

Lemma 5.6 *The number of vertices in category (2) (i.e., the number of vertices not in $V[\mathfrak{R}]$) is bounded by $18k$.*

PROOF. Let v and w be any two distinct vertices in D and observe the following. First, $N^{\dagger}(v) \cap N^{\dagger}(w) = \emptyset$, because if $u \in N^{\dagger}(v) \cap N^{\dagger}(w)$ then (v, u, w) would be a degenerated region with $u \notin V[\mathfrak{R}]$ contradicting the maximality of \mathfrak{R} . Second, from the first observation it follows that $w \notin N_{\dagger}^*(v)$ and $v \notin N_{\dagger}^*(w)$, and in general no vertex $a \in D$ belongs to $N_{\dagger}^*(b)$ for any vertex $b \in D$; otherwise, there exists a vertex $u \in N^{\dagger}(v)$ that is connected to w , and hence $u \in N^{\dagger}(v) \cap N^{\dagger}(w)$, contradicting the first observation. Third, $N_{\dagger}^*(v) \cap N_{\dagger}^*(w) = \emptyset$; otherwise, there exists a vertex $u \in N_{\dagger}^*(v) \cap N_{\dagger}^*(w)$ that is connected to a category-(2) vertex $a \in N^{\dagger}(v)$ (or $b \in N^{\dagger}(w)$) and the degenerated region (v, a, u, w) (or (w, b, u, v)) would contain the vertex $a \notin \mathfrak{R}$ (or $b \notin \mathfrak{R}$), contradicting the maximality of \mathfrak{R} .

Let B be the number of vertices not in D that are boundary-adjacent to vertices in D (i.e., in $N^*(v) - D$ for some $v \in D$). Combining the above observations with Proposition 5.5, it follows that the number of category-(2) vertices is

$$\sum_{v \in D} |N^{\dagger}(v)| \leq \frac{3}{2} \sum_{v \in D} |N_{\dagger}^*(v)| \leq 3B/2.$$

According to the definition of a region, each region in \mathfrak{R} has at most six vertices on its boundary two of which are vertices in D . Thus, each region in \mathfrak{R} can contribute with at most four vertices to

B . Note that from the above discussion no vertex $a \in D$ belongs to $N_{\dagger}^*(b)$ for any vertex $b \in D$, and hence the endpoints of the regions do not contribute to B . By Corollary 5.2, the number of regions in \mathfrak{R} is bounded by $3k - 6$. It follows that $B \leq 12k - 24$, and hence, the number of category-(2) vertices is bounded by $18k - 36 < 18k$. This completes the proof. \square

To bound the number of vertices in category (1), fix a region $R(v, w)$ between $v, w \in D$. We have the following lemma whose proof is technical and based on case-by-case structural analysis. The proof of the lemma appears in the appendix.

Lemma 5.7 (Appendix, Lemma 7.6) *Let $R = R(v, w)$ be a region in $V[\mathfrak{R}]$. The number of vertices in $V(R)$ is bounded by 16.*

Theorem 5.8 *The number of vertices in G is bounded by $67k$.*

PROOF. By Lemma 5.6, the number of category-(2) vertices in G is bounded by $18k$. Using this bound, we can assume that each region in \mathfrak{R} is nice. By Corollary 5.2, the number of regions in \mathfrak{R} is bounded by $3k - 6$. According to Lemma 5.7, the number of vertices in $V(R)$, where $R \in \mathfrak{R}$ is a nice region, is bounded by 16. It follows that the number of vertices in $V(\mathfrak{R})$ is bounded by $48k - 96$. Thus, the number of vertices in $V[\mathfrak{R}]$, and hence, in category (1), is bounded by $48k - 96$ plus the number of vertices in D which are the endpoints of the regions in \mathfrak{R} . Therefore the number of vertices in $V[\mathfrak{R}]$ is bounded by $49k - 96$, and the total number of vertices in G is bounded by $67k - 96 < 67k$. This completes the proof. \square

Theorem 5.9 *Let G be a planar graph with n vertices. Then in time $O(n^3)$, computing a dominating set for G of size bounded by k can be reduced to computing a dominating set of size bounded by k , for a planar graph G' of $n' < n$ vertices, where $n' \leq 67k$.*

PROOF. According to Theorem 4.1, in time $O(n^3)$ we can construct a reduced graph G' from G where $\gamma(G') = \gamma(G)$, and such that a dominating set for G can be constructed from a dominating set for G' in linear time. Moreover, the graph G' has no more than n vertices. If G has a dominating set of size bounded by k , then G' has a dominating set of size bounded by k (since $\gamma(G) = \gamma(G')$), and by Theorem 5.9, we must have $n' \leq 67k$. If this is the case, then we can work on computing a dominating set for G' of size bounded by k , from which a dominating set for G can be easily computed. If this is not the case, then G does not have a dominating set of size bounded by k , and the answer to the input instance is negative. This completes the proof. \square

6 A simple algorithm

In this section we present a simple algorithm for determining whether a graph G has a dominating set of size bounded by k .

Let $G = (V, E)$ be a planar graph given with an embedding in the plane. The *layer decomposition* of G with respect to the embedding, is a partitioning of V into disjoint layers (L_1, \dots, L_r) defined inductively as follows. Layer L_1 is the set of vertices that lie on the outer face of G , and layer L_i is the set of vertices that lie on the outer face of $G - \bigcup_{j=1}^{i-1} L_j$ for $1 < i \leq r$. It is well-known that a layer decomposition of a planar graph G can be computed in linear time in the number of vertices in the graph [4].

A *separator* in a graph G is a set of vertices S whose removal disconnects G . If (L_1, \dots, L_r) is a layer decomposition of G , then clearly the vertices in any layer L_i form a separator in G , separating

the vertices in layers L_1, \dots, L_{i-1} from those in layers L_{i+1}, \dots, L_r . Let (G, k) be an instance of the PLANAR-DS problem. By Theorem 5.9, we can assume that G is reduced, and that the number of vertices n of G satisfies $n \leq 67k$. Let (L_1, \dots, L_r) be a layer decomposition of G . Let $c > 0$ be a constant which will be determined later, and set $l = \lceil c\sqrt{k} \rceil$. Consider the families of layers F_i , $i = 1, \dots, l$, where F_i consists of layers $L_i, L_{i+l}, L_{i+2l}, \dots$. Assume for now that the number of layers $r \geq l$. We will show later how to handle the situation when this is not the case. The families F_i , $i = 1, \dots, l$, are disjoint, and each family forms a separator separating the graph into connected components that will be called *chunks*, where each chunk consists of at most l consecutive layers. Since these l families are disjoint and partition the layers into l groups, and since the graph has at most $67k$ vertices, there exists an index $1 \leq \mu \leq l$, such that the number of vertices in F_μ is bounded by $67k/l$. Again, observe that the removal of F_μ from G separates G into chunks, each consisting of at most l consecutive layers. Let these chunks be G_1, \dots, G_s .

The basic idea behind the algorithm is to apply a simple divide-and-conquer strategy by removing the vertices in the family F_μ to split the graph into chunks, then to compute a minimum dominating set for the resulting chunks using the algorithm introduced in [33], which is a variation of Baker’s algorithm [8]. To do this, for each vertex v in the F_μ , we “guess” whether v is in the minimum dominating set for G or not (basically, what we mean by guessing is enumerating all sequences corresponding to the different possibilities). For each guess of all the vertices in F_μ , we will solve the corresponding instance with respect to that guess. It was shown in [33] how this guessing process can be achieved using at most three statuses per vertex. Hence, guessing the vertices in F_μ can be done by enumerating at most $3^{|F_\mu|} \leq 3^{67k/l}$ ternary sequences. After guessing each vertex in the separator and updating the graph accordingly, the instance becomes an instance of a variation of the minimum dominating set problem due to the constraints placed on some of the vertices in the graph. Kanj and Perković introduced an algorithm in [33], which is a variation of Baker’s algorithm [8], to solve this problem. The algorithm introduced in [33] solves this problem on the chunks in time $O(27^{d+1}n)$, where d is the maximum number of layers in a chunk (i.e., the maximum depth of a chunk). Noting that $d \leq l$ and that $n \leq 67k$, we conclude that after guessing all the vertices in F_μ , the problem can be solved in time $O(27^l k)$. If the number of layers r in G is less than l , we can simply call the algorithm in [33] directly on G to solve the problem in time $O(27^l k)$. The algorithm is given in Figure 3 below.

It is not difficult to verify that the running time of the algorithm is $O(3^{67k/l} \cdot 27^l \cdot k + n^3)$, where the $O(n^3)$ time is to count for the time taken to reduce G to its kernel. Niedermeier and Rossmanith showed how to get rid of the k factor corresponding to the kernel size in the running time of such algorithms [36]. Using their techniques, the running time of the algorithm becomes $O(n^3 + 3^{67k/l} \cdot 27^l)$. We choose c , and hence l , so that the above expression is minimized. It can be shown that the expression is minimized when $c = \sqrt{67/3}$, and the running time of the algorithm becomes $O(n^3 + 2^{45\sqrt{k}})$.

Theorem 6.1 *In time $O(n^3 + 2^{45\sqrt{k}})$, it can be determined whether a planar graph on n vertices has a dominating set of size bounded by k or not.*

Theorem 6.1 shows that our algorithm for solving the PLANAR-DS problem is competitive with the previous algorithms using the similar technique of layer decomposition of a planar graph [4, 33]. The above algorithm improves the original $O(2^{70\sqrt{k}}n)$ time algorithm given in [4] for the problem. At the same time, our algorithm is much simpler than the algorithms in [4, 23, 25, 33], illustrating the power of kernelization in the process of designing efficient algorithms for parameterized \mathcal{NP} -hard problems. Finally, after a kind of race resulting in better and better algorithms [4, 23, 25, 33],

ALGORITHM. DS-solverInput: a planar graph G of n vertices, and a parameter k Output: a dominating set D of size $\leq k$ in case it exists;

1. use the results in Theorem 5.9 to kernelize G ;
2. **if** the number of vertices n of G is $> 67k$ **then**
 Stop(“ G does not have a dominating set of size $\leq k$ ”);
3. let $c = \sqrt{67/3}$; $l = \lceil c\sqrt{k} \rceil$;
4. **if** the number of layers in G is $< l$ **then**
 use the algorithm in [33] to solve the problem in time $O(2^{45\sqrt{k}}k)$; **Stop**;
5. let F_μ be a separator of size $\leq 67k/l$ separating the graph into chunks G_1, \dots, G_s
 each consisting of at most l consecutive layers;
6. **for** each assignment to the vertices in F_μ **do**
 update D ;
 split the graph into its components;
 compute a minimum dominating set D' for the resulting graph using the algorithm in [33];
 $D = D \cup D'$;
7. output the smallest dominating set constructed in step 6 in case its size is bounded by k ; otherwise
 return (“ G does not have a dominating set of size $\leq k$ ”).

Figure 3: A simple algorithm solving PLANAR-DS.

recently Fomin and Thilikos presented an $O(n^3 + 2^{15.13\sqrt{k}})$ time algorithm to solve the PLANAR-DS problem based on the concept of branchwidth [25]; the best treewidth based algorithm being only slightly worse [23].

7 Summary and Extensions

In this paper we exhibited the first lower bound results on kernel sizes and, motivated by this findings, we strived to improve on the (still huge) constants involved in the known linear kernel for PLANAR-DS.

Are there other, possibly more sophisticated arguments for showing lower bounds on kernel sizes? In particular, it would be interesting to have arguments ruling out, for example, the existence of a kernel of size $o(k^3)$ in a situation when a kernel of size $\mathcal{O}(k^3)$ has been obtained. The algebra we used in the proof of Theorem 3.1 does not extend to such cases.

We mention that the concept of a black-and-white graph we used for deriving the kernel upper bound results for PLANAR-DS also allows us to exhibit a small kernel for a variation of the DOMINATING SET problem, called the RED/BLUE DOMINATING SET problem, as studied by Weihe [41]: Given a graph $G = (V, E)$ with V partitioned into $V_{\text{red}} \cup V_{\text{blue}}$, and a positive integer k , is there a *red/blue dominating set* $D \subseteq V_{\text{red}}$ with $|D| \leq k$, i.e., $V_{\text{blue}} \subseteq N(D)$? Namely, if we consider the red vertices as “black” in our notation and the blue vertices as “white”, and if we re-analyze our reduction rules, we can state:

Corollary 7.1 PLANAR RED/BLUE DOMINATING SET *admits a problem kernel of size $67k$.*

As exhibited in [4, 23], the possibly better known FACE COVER problem can be solved with the help of PLANAR RED/BLUE DOMINATING SET, by introducing “face vertices.” However, we are still investigating if we could claim a small kernel for FACE COVER, since we are not keeping the (face) structure of the original problem. Notice that a cubic kernel was derived in [1, Theorem 1]. Based on this sort of problem kernel, we could then arrive at an $O^*(c^{\sqrt{k}})$ algorithm for FACE COVER that

is significantly better than what was obtained in [4], close to being competitive with [23], along the lines of the preceding section.

References

- [1] F. ABU-KHZAM AND M. LANGSTON, A direct algorithm for the parameterized face cover problem, in *Proceedings of the 1st International Workshop on Parameterized and Exact Computation, Lecture Notes in Computer Science* **3162**, (2004), pp. 213–222.
- [2] F. ABU-KHZAM, M. LANGSTON, AND P. SHANBHAG, A High-Performance Toolkit for Fast Exact Algorithms, invited talk to the *Workshop on Fixed-Parameter Tractability*, Ottawa, Canada, 2003. Available at <http://citeseer.nj.nec.com/560785.html>.
- [3] J. ALBER, N. BETZLER, AND R. NIEDERMEIER, Experiments on data reduction for optimal domination in networks, *Annals of Operations Research* **146**, (2006), pp. 105–117.
- [4] J. ALBER, H. BODLAENDER, H. FERNAU, T. KLOKS, AND R. NIEDERMEIER, Fixed parameter algorithms for Dominating Set and related problems on planar graphs, *Algorithmica* **33**, (2002), pp. 461–493.
- [5] J. ALBER, M. FELLOWS, AND R. NIEDERMEIER, Polynomial-time data reduction for dominating set, *Journal of the ACM* **51**, (2004), pp. 363–384.
- [6] J. ALBER, H. FERNAU, AND R. NIEDERMEIER, Parameterized complexity: exponential speedup for planar graph problems, *Journal of Algorithms* **52**, (2004), pp. 26–56.
- [7] J. ALBER, H. FERNAU, AND R. NIEDERMEIER, Graph separators: a parameterized view, *Journal of Computer and System Sciences* **67**, (2003), pp. 808–832.
- [8] B. BAKER, Approximation algorithms for NP-complete problems on planar graphs, *Journal of the ACM* **41**, (1994), pp. 153–180.
- [9] R. BALASUBRAMANIAN, M. R. FELLOWS, AND V. RAMAN, An improved fixed parameter algorithm for vertex cover, *Information Processing Letters* **65**, (1998), pp. 163–168.
- [10] J. CHEETHAM, F. DEHNE, A. RAU-CHAPLIN, U. STEGE, AND P. J. TAILLON, Solving large FPT problems on coarse grained parallel machines, *Journal of Computer and System Sciences* **67**, (2003), pp. 691–706.
- [11] J. CHEN AND I. KANJ, Constrained Minimum Vertex Cover in Bipartite Graphs: Complexity and Parameterized Algorithms, *Journal of Computer and System Sciences* **67**, (2003), pp. 833–847.
- [12] J. CHEN AND I. KANJ, Improved Exact Algorithms for MAX-SAT, *Discrete Applied Mathematics* **142(1-3)**, (2004), pp. 17–27.
- [13] J. CHEN, I. KANJ, AND W. JIA, Vertex cover: further observations and further improvement, *Journal of Algorithms* **41**, (2001), pp. 280–301.
- [14] J. CHEN, I. KANJ, AND G. XIA, Improved parameterized upper bounds for vertex cover, in *Proceedings of the 31 International Symposium on the Mathematical Foundations of Computer Science, Lecture Notes in Computer Science* **4162**, (2006), pp. 238–249.
- [15] F. Dehne, M. Fellows, H. Fernau, E. Prieto, and F. Rosamond. NONBLOCKER: parameterized algorithmics for MINIMUM DOMINATING SET, in *Software Seminar SOFSEM, Lecture Notes in Computer Science* **3831**, (2006), pp. 137–147.

- [16] R. DIESTEL, *Graph Theory*, Springer, (1996).
- [17] R. DOWNEY AND M. FELLOWS, *Parameterized Complexity*, Springer, (1999).
- [18] R. DOWNEY, M. FELLOWS, AND U. STEGE, Parameterized Complexity: A Framework for Systematically Confronting Computational Intractability, in *Contemporary Trends in Discrete Mathematics*, (R. Graham, J. Kratochvíl, J. Nešetřil, and F. Roberts eds.), proceedings of the DIMACS-DIMATIA Workshop, Prague 1997, *AMS-DIMACS Series in Discrete Mathematics and Theoretical Computer Science* vol. **49**, (1999), pp. 49–99.
- [19] M. FELLOWS, Parameterized complexity: the main ideas and connections to practical computing, in *Electronic Notes in Theoretical Computer Science* **61**, (2002).
- [20] M. FELLOWS, C. MCCARTIN, F. ROSAMOND, AND U. STEGE, Coordinatized kernels and catalytic reductions: an improved FPT algorithm for Max Leaf Spanning Tree and other problems, in *FST TCS 2000, Lecture Notes in Computer Science* **1974**, (2000), pp. 240–251.
- [21] H. FERNAU, On parameterized enumeration, in *Computing and Combinatorics, Proceedings COCOON, Lecture Notes in Computer Science* **2383**, (2002), pp. 564–573.
- [22] H. FERNAU, ROMAN DOMINATION: a parameterized perspective, in *Software Seminar SOFSEM*, (2006), *Lecture Notes in Computer Science* **3831**, (2006), pp. 262–271.
- [23] H. FERNAU AND D. JUEDES, A geometric approach to parameterized algorithms for domination problems on planar graphs, in *Mathematical Foundations of Computer Science MFCS 2004, Lecture Notes in Computer Science* **3153**, (2004), pp. 488–499.
- [24] H. FERNAU AND R. NIEDERMEIER, An efficient exact algorithm for constraint bipartite vertex cover. *Journal of Algorithms* **38**, (2001), pp. 374–410.
- [25] F. FOMIN AND D. THILIKOS, Dominating sets in planar graphs: branch-width and exponential speed-up, in *Proceedings of the 14th ACM-SIAM Symposium on Discrete Algorithms SODA*, (2003), pp. 168–177.
- [26] F. V. FOMIN AND D. THILIKOS, Fast parameterized algorithms for graphs on surfaces: Linear kernel and exponential speed-up, in *International Colloquium on Automata, Languages and Programming ICALP, Lecture Notes in Computer Science* **3142**, (2004), pp. 581–592.
- [27] M. GAREY AND D. JOHNSON, *Computers and Intractability: A Guide to the Theory of NP-completeness*, Freeman, San Francisco, 1979.
- [28] J. GIMBEL AND C. THOMASSEN, Coloring triangle-free graphs with fixed size, *Discrete Mathematics* **219**, (2000), pp. 275–277.
- [29] H. GRÖTZSCH, Ein Dreifarbensatz für dreikreisfreie Netze auf der Kugel, *Wiss. Zeitschrift der Martin-Luther-Univ. Halle-Wittenberg, Math.-Naturwiss. Reihe* **8**, (1959), pp. 109–120.
- [30] F. HENGLEIN AND H. G. MAIRSON, The complexity of type inference for higher order typed lambda calculi, *Journal of Functional Programming* **4**, (1994), pp. 435–477.
- [31] J. HOPCROFT AND R. TARJAN, Efficient planarity testing, *Journal of the ACM* **21**, (1974), pp. 549–568.
- [32] R. HYAFIL AND R. L. RIVEST, Constructing optimal binary trees is NP-complete, *Information Processing Letters* **5**, (1976), pp. 15–17.

- [33] I. KANJ AND L. PERKOVIĆ, Improved parameterized algorithms for planar dominating set, in *Mathematical Foundations of Computer Science MFCS, Lecture Notes in Computer Science* **2420**, (2002), pp. 399–410.
- [34] S. KHOT AND V. RAMAN, Parameterized complexity of finding subgraphs with hereditary properties, *Theoretical Computer Science* **289**, (2002), pp. 997–1008.
- [35] O. LICHTENSTEIN AND A. PNEULI, Checking that finite-state concurrents programs satisfy their linear specification, *Proc. 12th ACM Symp. on Principles of Programming Languages POPL*, (1985), pp. 97–107.
- [36] R. NIEDERMEIER AND P. ROSSMANITH, A general method to speed up fixed-parameter-tractable algorithms, *Information Processing Letters* **73**, (2000), pp. 125–129.
- [37] O. ORE, *Theory of Graphs*, Colloquium Publications XXXVIII, *American Mathematical Society*, (1962).
- [38] E. PRIETO AND C. SLOPER, Either/or: Using vertex cover structure in designing FPT-algorithms-the case of k -internal spanning tree, in *Proceedings Workshop on Algorithms and Data Structures WADS, Lecture Notes in Computer Science* **2748**, (2003), pp. 465–483.
- [39] C. THOMASSEN, Grötzsch’s 3-color theorem and its counterparts for the torus and the projective plane, *Journal of Combinatorial Theory, Series B* **62**, (1994), pp. 268–279.
- [40] R. UEHARA, *Probabilistic Algorithms and Complexity Classes*, PhD thesis, Department of Computer Science and Information Mathematics, The University of Electro-Communications, Japan, March 1998.
- [41] K. WEIHE, Covering trains by stations or the power of data reduction, in *Algorithms and Experiments ALEX*, (1998), pp. 1–8. <http://rtm.science.unitn.it/alex98/proceedings.html>.

Appendix

In this section we prove Lemma 5.7. We first start with some observations and preliminary results.

Fix a region $R(v, w)$ between $v, w \in D$. Without loss of generality, assume the boundary of R is determined by the two paths (v, v_1, w_1, w) and (v, v_2, w_2, w) . Note that all vertices in $V(R)$ belong to $N(v, w)$, and that $v_1, v_2 \in N^*(v)$, and $w_1, w_2 \in N^*(w)$. If there is a degree-1 vertex x connected to v (resp. w), then this vertex is in $N_3(v)$ (resp. $N_3(w)$) and must be colored white by **Rule 3**. Similarly, if there exists a degree-2 vertex y that is connected to v and either v_1 or v_2 (resp. w and either w_1 or w_2), then y is in $N_2(v)$ (resp. $N_2(w)$) and must be colored white by **Rule 3**. Now if a degree-1 white vertex is connected to v , then since the vertices in $N^\dagger(v)$ are white and are neighbors of v , by **Rule 6** we must have $N^\dagger(v) = \emptyset$. During the process of counting the number of vertices in $N^\dagger(v)$, we bounded the number of vertices in $N^\dagger(v)$ by $3|N^*(v)|/2$. This can be looked at as each vertex in $N^*(v)$ contributing $3/2$ vertices to $|N^\dagger(v)|$. So if a degree-1 white vertex is connected to v (note that at most one degree-1 vertex can be connected to v), this means that $N^*(v)$ which contains at least two vertices will not contribute to the number of vertices in $N^\dagger(v)$, and hence, the bound on $|N^\dagger(v)|$ will be decreased by at least three. Similarly, if a degree-2 white vertex is connected to v and v_1 (or v_2) (again, note that there can be at most one degree-2 vertex connected to v and to v_1 (or v_2)), then no vertex in $N^\dagger(v)$ can be connected to v_1 (or v_2). This can be regarded as a reduction to the bound on $|N^\dagger(v)|$ by $3/2$. Thus, if we use the upper bound on the number of vertices in category (2) computed above, we may assume without loss of generality that no degree-1 vertex is connected to v or w , and that no degree-2 vertex is connected to v and v_1, v and v_2, w and w_1, w and w_2 . We will also assume that the boundary of a region $R(v, w)$ consists of exactly six distinct vertices, that is, the region is not a degenerate region. The case of a degenerate region obviously yields a better bound on the number of vertices in the region. Let us call a region with all the above properties *nice*. We start with the following propositions.

Proposition 7.2 *If there is no simple black vertex strictly inside a quasi-simple region $R = R(v, w)$, then $V(R)$ contains at most two simple white vertices.*

PROOF. Suppose, to get a contradiction, that $V(R)$ contains more than two simple white vertices, and let a, b , and c be three such vertices. Since all the three vertices are simple, one vertex must be engulfed within the area determined by v, w , and the other two vertices. Suppose that b is situated within the area (v, a, w, c, v) . Since, by the assumption of the proposition, all the simple vertices strictly inside $V(R)$ must be white, and since all the non-simple vertices inside $V(R)$ (i.e., vertices in R^+) are white by definition, and since no edges exist between white vertices, it follows that the white simple vertex b , engulfed by the area (v, a, w, c, v) , is only connected to v and w , and hence has degree 2. Note that the color of both v and w must be black since there are simple white vertices that are connected to both v and w . Now $|W(v, w)| > 2$ because $\{a, b, c\} \subseteq W(v, w)$. But this makes **Rule 8** applicable contradicting the fact that G is reduced. This completes the proof. \square

Proposition 7.3 *Let $R = R(v, w)$ be a quasi-simple region where the color of v is black, then $V(R)$ has at most 4 simple vertices.*

PROOF. Suppose first that R has six or more simple vertices. Let S be the set of those simple vertices that are strictly inside R . Then $|S| \geq 4$. Since the vertices in S are simple and hence connect to both v and w , it is obvious that no vertex lying strictly inside R can dominate all vertices in S . But S is a subset of those vertices in $N_2(v, w) \cup N_3(v, w)$ that are strictly inside R , it

follows that no vertex that is strictly inside R can dominate all vertices in $N_2(v, w) \cup N_3(v, w)$. Now all vertices that lie strictly inside R belong to $N_2(v, w) \cup N_3(v, w)$, thus, by **Rule 5**, all vertices strictly inside R must be white. Noting that $|S| \geq 4$, and that all the vertices in S are simple white vertices, this contradicts Proposition 7.2.

Suppose now that R has five simple vertices. Let a , b , and c , be the three simple vertices that lie strictly inside R . By an argument similar to the above, we can assume that vertex b is engulfed within the area determined by v , a , w , and c . Again all the vertices strictly inside R must belong to $N_2(v, w) \cup N_3(v, w)$. Since a does not dominate c , and vice versa, it follows that a and c are colored white by **Rule 5**. Now a , b , and c are the only simple vertices strictly inside R , by Proposition 7.2, no three simple white vertices can be contained in R . This forces b to be black, and to be connected to both a and c (otherwise b would be colored white by **Rule 5**). Now all other non-simple vertices in R must be connected to the boundary, and hence cannot be connected to b (all the vertices other than a and c which can be connected to b have to belong to the area engulfed by (v, a, w, c) and cannot be connected to the boundary). Thus, $W(b) = \{a, c\}$, and every vertex in $W(b)$ is connected to all vertices in $B(b) = \{v, w\}$ (note that since a and c are white, and are connected to w , w must be black). By **Rule 7**, $W(b) = \{a, c\}$ should have been removed at this point, a contradiction. Therefore, R has at most four simple vertices and the proof is complete. \square

Proposition 7.4 *Let (v, y, w, z, v) be the boundary of a quasi-simple region $R = R(v, w)$, and suppose that v and y are black. Then there can be at most one white vertex in $R^+ = R^+(v, w)$ that is connected to both v and y .*

PROOF. Suppose, to get a contradiction, that there are at least two white vertices in R^+ that are connected to both v and y . Since all the vertices in R^+ are white, and hence cannot be connected together, there must exist two white vertices a and b in R^+ satisfying that the area engulfed by (v, a, y, v) is empty, and the area engulfed by (v, b, y, v) contains only the vertex a . Clearly, the degree of a is exactly 2, and a belongs to $N_2(v) \cup N_3(v)$. Now since both a and b are connected to both v and y , we have $N(a) \subseteq N(b)$. Given the fact that v is black, this is a contradiction to **Rule 6**. \square

Proposition 7.5 *Let $R = R(v, w)$ be a quasi-simple region, and suppose that v is black. Let (v, y, w, z, v) be the boundary of R . If*

- (a) *there are no simple vertices strictly inside R , or*
- (b) *there are simple vertices strictly inside R and all vertices in $R^+ = R^+(v, w)$ are connected to y ,*

then $V(R) \cup \{w\}$ contains at most three white vertices. Moreover, if there are three white vertices in $V(R) \cup \{w\}$, then either $R^+ \neq \emptyset$, or there is a simple black vertex interior to R .

PROOF. To prove that part (a) implies the statement of the proposition, suppose that there are no simple vertices lying strictly inside R . Then clearly all the white vertices in $V(R)$ come from $R^+ \cup \{y, z\}$. If y is white, then no vertex in R^+ can be connected to y because the vertices in R^+ are all white. On the other hand, since $R^+ \subseteq N_2(v) \cup N_3(v)$, if y is black, by Proposition 7.4, at most one white vertex in R^+ can be connected to y . Similarly if z is black. Since every vertex in R^+ has to be connected to either y or z by the definition of a quasi-simple region, it follows from the above that $V(R)$ contains at most two white vertices, and hence $V(R) \cup \{w\}$ contains at most

three white vertices. Now when $V(R) \cup \{w\}$ contains three white vertices, w must be white, and hence, y and z are black. Thus, the two white vertices other than w in $V(R) \cup \{w\}$ come from R^+ , and $R^+ \neq \emptyset$.

To prove part (b), note first that, by Proposition 7.3, the number of simple vertices in R including y and z is bounded by four. We will assume that the number of simple vertices in R is exactly four. The cases when there are less than four simple vertices in R are simpler, and yield the desired bound. Let a and b be the other two simple vertices, and assume that the four simple vertices y, a, b, z appear in the preceding sequence in a clockwise order around v . Observe that the white vertices in $V(R)$ come from $R^+ \cup \{y, a, b, z\}$. Also observe that since all the vertices in R^+ are connected to y by the hypothesis of part (b), either y is white and R^+ is empty, or y is black and by Proposition 7.4, R^+ contains at most one vertex. It follows that the number of white vertices in $R^+ \cup \{y\}$ is bounded by one. Now suppose to get a contradiction that $V(R) \cup \{w\}$ contains four white vertices. Since no two white vertices are connected, and since all vertices in $\{a, b, z\}$ are connected to w , w must be black and all the three vertices a, b , and z must be white. But then the degree of b is exactly 2, and $|W(v, w)| > 2$, contradicting **Rule 8**. To complete the proof, suppose that $V(R) \cup \{w\}$ contains exactly three white vertices, we need to show that either $R^+ \neq \emptyset$ or there exists a simple black vertex inside R . Suppose to get a contradiction that $R^+ = \emptyset$ and the interior vertices to R , a, b , are all white. Then w must be black in this case and either y or z is white. Without loss of generality, assume y is white. Since there are no edges between white vertices, the degree of a must be two and $\{y, a, b\} \subseteq W(v, w)$, again a contradiction to **Rule 8**. \square

Lemma 7.6 (Lemma 5.7) *Let $R = R(v, w)$ be a nice region in $V[\mathfrak{R}]$. The number of vertices in $V(R)$ is bounded by 16.*

PROOF. Every vertex in $V(R)$ is in $N(v, w)$, and hence, is either connected to v or w . We distinguish two cases.

Case 1. $N_3(v, w) = \emptyset$. In this case every vertex in $V(R) - \{v_1, v_2, w_1, w_2\}$ has to be connected to at least one vertex in $\{v_1, v_2, w_1, w_2\}$ because v_1, v_2, w_1, w_2 are the only vertices in $V(R)$ that possibly belong to $N_1(v, w)$. Since R is nice, the vertices in $V(R) - \{v_1, v_2, w_1, w_2\}$ can be classified into the following categories, where a vertex is assigned to the first category that it satisfies:

- (i) vertices connected to v and v_1 , but not connected to w_1 ;
- (ii) vertices connected to v and w_1 ;
- (iii) vertices connected to v and w_2 ;
- (iv) vertices connected to v and v_2 ;
- (v) vertices connected to w and w_2 , but not connected to v_2 ;
- (vi) vertices connected to w and v_2 ;
- (vii) vertices connected to w and v_1 ; and
- (viii) vertices connected to w and w_1 .

Note that one of categories (ii) and (vii) must be empty, otherwise, according to our placement of the vertices in the categories, we have two distinct vertices in $V(R)$ other than v_1 and w_1 , one of them is connected to v and w_1 and the other to w and v_1 , contradicting the planarity of the graph. Similarly, one of categories (iii) and (vi) must be empty. Without loss of generality, assume that categories (iii) and (vii) are empty. If, in addition, any of categories (ii) or (vi) is empty, then the situation becomes simpler leading to a better bound on the number of vertices in $V(R)$. Thus,




Figure 4: Illustration of a possible worst-case scenario for **Case 1**. Empty circles represent white vertices, and filled circles black vertices.

we will assume that both categories (ii) and (vi) are nonempty. Note also that since R is nice, no vertex interior to R has degree 2, and every vertex in category (i) must be connected to some vertex interior to R . Since category (ii) is nonempty, and by the planarity of G , a vertex in category (i) can only be connected to vertices in category (ii). Moreover, since the vertices in category (i) are only connected to v , and to neighbors of v including v_1 (since these vertices can only be connected to v and to vertices in category (ii)), all these vertices belong to $N_2(v) \cup N_3(v)$. Since v is black, by **Rule 3**, all vertices in category (i) must be white. Now the vertices in category (i) and category (ii), plus v , v_1 , and w_1 , form a quasi-simple region between v and w_1 , $Q_1 = R(v, w_1)$. Moreover, all the vertices in $V(Q_1)$, except those in category (i), are simple vertices because all vertices in $V(Q_1)$, except those in category (i), have to be connected to both v and w_1 . Since v is black (all the endpoints of regions in \mathfrak{R} are black), by Proposition 7.3, the number of vertices in $V(Q_1)$ except those vertices in category (i), is bounded by 4. Now we bound the number of vertices in category (i). Every vertex in category (i) is white and is connected to v and v_1 . If category (i) is nonempty, then v_1 must be black, and the vertices in category (i) are white vertices in $Q_1^+(v, w_1)$ that are connected to v and v_1 . It follows from Proposition 7.4 that the number of vertices in category (i) is bounded by 1. This shows that the number of vertices in $V(Q_1)$ is bounded by five. By symmetry, the number of vertices in $V(Q_2)$, where Q_2 is the quasi-simple region between w and v_2 consisting of the vertices in category (v) and category (vi), plus the vertices w , w_2 , v_2 , is bounded by five. Now we bound the number of vertices in categories (iv) and (viii). We have the following claim.

Claim. The number of vertices in category (iv) (resp. category (viii)) is bounded by 2. Moreover, at most one vertex in category (iv) (resp. category (viii)) is white.

Consider the vertices in category (iv). Suppose that there are three or more vertices in category (iv), and let a_1, a_2, a_3 be three vertices in category (iv) such that: no vertex is engulfed in the area of the embedding determined by (v, a_1, v_2) , a_1 is the only vertex engulfed in the area determined by (v, a_2, v_2) , and a_1 and a_2 are the only two vertices engulfed in the area determined by (v, a_3, v_2) . Now a_1 and a_2 must belong to $N_2(v) \cup N_3(v)$. Since v is black, it follows from **Rule 3** that a_1 and

a_2 must be white, and no edge exists between a_1 and a_2 . But this means that $N(a_1) \subseteq N(a_2)$, and since $a_1 \in N_2(v)$ and v is black, then according to **Rule 6**, this leads to a contradiction. It follows that at most two vertices can be in category (iv). By symmetry, at most two vertices can be in category (viii). Note also that it follows from the above proof that if there are exactly two vertices a_1 and a_2 in category (iv) (resp. category (viii)), then at most one vertex in $\{a_1, a_2\}$ can be white. This proves the claim.

Now the vertices in $V(R)$ consist of vertices of $V(Q_1)$, vertices of $V(Q_2)$, category (iv) and category (viii) vertices, and the two vertices v_2 and w_1 (in case these two vertices were not included in any of the other categories). It follows that the number of vertices in R is bounded by 16. See Figure 4 for an illustration of such a possible scenario.

Case 2. $N_3(v, w) \neq \emptyset$. Let X be the set of white vertices in $N_2(v, w)$ that are in $V(R)$, Y the set of black vertices in $N_2(v, w) \cup N_3(v, w)$ that are in $V(R)$, and Z the set of white vertices in $N_3(v, w)$ that are in $V(R)$. We first draw few observations.

Observation 1. $|X| \leq 7$. We first show that $|X| \leq 8$. Remove the vertices in $N_3(v, w)$ interior to R , then define categories (i) – (viii) as above. Similar to **Case 1**, we can assume that the vertices in category (i) and (ii), plus the vertices v, v_1 , and w_1 , form a quasi-simple region $Q_1 = R(v, w_1)$, and those in category (v) and (vi), plus the vertices w, w_2 , and v_2 , form a quasi-simple region $Q_2 = R(w, v_2)$. Since $X \subseteq N_2(v, w)$, every vertex in X must belong to one of categories (i) – (viii), or possibly to the set $\{v_1, v_2, w_1, w_2\}$. From the definition of categories (i) and (ii), vertices in category (i) form the set $Q_1^+ = Q_1^+(v, w_1)$ in the quasi-simple region Q_1 , and all the vertices in Q_1^+ are connected to v_1 . Now add the vertices in $N_3(v, w)$ back, and note that no black vertex in $N_3(v, w)$ that is not connected to w_1 resides in Q_1 . The reason being that such a vertex would be in $N_2(v) \cup N_3(v)$ (otherwise, this vertex will have to be connected to w_1 — the only vertex in Q_1 possibly not in $N(v)$) and hence colored white by **Rule 3**. Now Q_1 plus the set of black vertices in $N_3(v, w)$ that reside in Q_1 , minus the set of white vertices in $N_3(v, w)$ that reside in Q_1 , satisfies condition (b) in Proposition 7.5, and the number of white vertices in Q_1 is bounded by three. Since no two white vertices are connected together, and hence the presence of the white vertices from $N_3(v, w)$ in Q_1 cannot increase the number of possible white vertices in Q_1 , we conclude that the number of white vertices in Q_1 that are not in $N_3(v, w)$, and hence, the number of vertices in X that belong to Q_1 is bounded by three. Similarly, the number of vertices in X that belong to Q_2 is bounded by three. Moreover, the statement of the claim in **Case 1** carries in a straightforward manner to **Case 2**, and categories (iv) and (viii) contain at most one white vertex each. It follows that the number of white vertices in the set X , is bounded by eight. Now if $|X| = 8$, then both Q_1 and Q_2 (plus the black vertices in $N_3(v, w)$ that reside in Q_1 and Q_2) contain three white vertices. Since Q_1 contains exactly three white vertices, by Proposition 7.5, either $Q_1^+ \neq \emptyset$, or Q_1 must contain an interior black vertex. If $Q_1^+ \neq \emptyset$, since R is nice, the vertex in Q_1^+ must be connected to some vertex interior to Q_1 which must be black because the vertices in Q_1^+ are white. Therefore, if Q_1 contains exactly three white vertices, then there must exist an interior black vertex p in Q_1 . Similarly, there must exist an interior black vertex q in Q_2 . Since both p and q are black and are in $N_2(v, w) \cup N_3(v, w)$, by **Rule 4**, p and q must dominate all vertices in $N_3(v, w) \neq \emptyset$. In particular, p which is interior to Q_1 must dominate q which is interior to Q_2 . This is a contradiction to the planarity of the graph. It follows that $|X| \leq 7$.

Observation 2. Every vertex in Y must dominate all vertices in $N_3(v, w)$.

This observation follows from **Rule 4** since the vertices in Y are black and are a subset of $N_2(v, w) \cup N_3(v, w)$.

Let H be the graph obtained from G by identifying the vertex v with w along the path (v, v_1, w_1, w) . Clearly, H is planar. Let u be the resulting vertex by this identification. Let Y' be the set of vertices in Y that are in H , and let $y = |Y'|$. Similarly, let Z' be the set of vertices in Z that are in H , and let $z = |Z'|$. Observe that the vertex u is connected to all the vertices in Y' and Z' in H , and that the only vertices that have been removed by this identification are boundary vertices to R that belong to $\{v_1, v_2, w_1, w_2\}$.

Observation 3. If $y > 1$ and $z > 1$, then the number of vertices in $V(R)$ is bounded by 16.

Suppose that $y > 1$ and $z > 1$. If $y > 2$, since every vertex in Y' must dominate the vertices in Z' , it follows that the subgraph of H induced by the set of vertices $Y' \cup Z' \cup \{u\}$ contains a copy of $K_{3,3}$, contradicting the planarity of H (the vertices in Y' form the first bipartition and the other vertices form the second bipartition). Suppose now that $y = 2$. If $z > 2$, then similarly, the subgraph induced by $Z' \cup \{u\} \cup Y'$ contains a copy of $K_{3,3}$ (the vertices in $Y' \cup \{u\}$ form the first bipartition, and those in Z' form the second bipartition). Suppose now that $y = z = 2$. Then the number of vertices in $X \cup Y' \cup Z'$ is bounded by 11. Since $|V(R)| \leq |X \cup Y' \cup Z' \cup \{v_1, v_2, w_1, w_2\}|$, it follows that the number of vertices in $V(R)$ is bounded by 16.

Now we distinguish the following two subcases.

Subcase 2.1. $z \leq 1$. Let $Y_1 = Y' \cap N_2(v, w)$ be the set of black vertices in Y' that are in $N_2(v, w)$, $y_1 = |Y_1|$, $Y_2 = Y' - Y_1$ be the set of black vertices in Y' that are in $N_3(v, w)$, and $y_2 = |Y_2|$. Note that every vertex in Y' must be connected to all vertices in $Y_2 \cup Z$ by **Rule 4**. If $y = y_1 + y_2 < 5$, then since $z \leq 1$ and the number of vertices in X is bounded by 7 by **Observation 1**, the number of vertices in $V(R)$ is bounded by 16. So we can assume that $y \geq 5$. If $y_2 + z \geq 4$, then the subgraph induced by the vertices $\{u\} \cup Y_2 \cup Z$ is a copy of K_5 . Thus, $y_2 + z < 4$. If $y_2 + z = 3$, then the subgraph induced by the bipartition $(Y_2 \cup Z, Y_1 \cup \{u\})$ contains a copy of $K_{3,3}$, whereas if $y_2 + z = 2$, then subgraph induced by the bipartition $(\{u\} \cup Y_2 \cup Z, Y_1)$ contains a copy of $K_{3,3}$. Suppose now that $y_2 + z = 1$. If $y_1 \leq 4$, then $y + z \leq 5$, and hence, the number of vertices in R is bounded by 16. Assume now that $y_1 \geq 5$. Let p be the vertex in $Y_2 \cup Z$, then p is connected to all vertices in Y_1 in H , and hence in G . Moreover, every vertex in Y_1 is either connected to v or w (or both) in G . Since $y_1 \geq 5$, there must exist at least three vertices in Y_1 that are connected either to v , or to w in G . Let these vertices be p_1, p_2 , and p_3 , and assume without loss of generality, that these vertices are connected to v . Since p_1, p_2 , and p_3 , are also connected to p , there must exist a vertex in $\{p_1, p_2, p_3\}$, say p_2 , that is interior to the region determined by v, p , and the other two vertices. But $p_2 \in Y_1 \subseteq N_2(v, w)$, and hence p_2 must be connected to the boundary of R (because p_2 must be connected to some vertex in $N_1(v, w)$), a contradiction. Thus, the number of vertices in $V(R)$ is bounded by 16.

Subcase 2.2. $y \leq 1$. If $z \leq 4$, then $y + z \leq 5$, and given that $|X| \leq 7$ by **Observation 1**, the total number of vertices in $V(R)$ is bounded by 16. Suppose now that $z \geq 5$. Observe first that $Y \neq \emptyset$, otherwise, $N_2(v, w) \cup N_3(v, w)$ consists only of white vertices, and $N_3(v, w)$, which contains at least five white vertices ($Z \subseteq N_3(v, w)$), could not be dominated by a single vertex

in $N_2(v, w) \cup N_3(v, w)$. This would make **Rule 2** applicable, a contradiction. Let $p_1, p_2, p_3, p_4,$ and p_5 be vertices in Z . Since each of these vertices must be connected to either w or v , at least three vertices in $\{p_1, p_2, p_3, p_4, p_5\}$ are connected either to v , or to w . Suppose, without loss of generality, that $\{p_1, p_2, p_3\}$ are connected to v , and note that by **Observation 2**, every vertex in Y must be connected to all vertices in Z . If $|Y| \geq 2$, then $(\{v\} \cup Y, \{p_1, p_2, p_3\})$ would be a copy of $K_{3,3}$. Suppose now that $|Y| = 1$, and let q be the single vertex in Y . Since p_1, p_2, p_3 are white and belong to $N_3(v, w)$, these vertices can only connect to vertices in $\{v, q, w\}$. Again, by planarity, at least one vertex in $\{p_1, p_2, p_3\}$, is not connected to w , and hence must be of degree 2. But then $|W(v, q)| \geq 3$, and $W(v, q)$ contains a degree-2 vertex. This contradicts **Rule 8**.

It follows that in all cases the number of vertices in $V(R)$ is bounded by 16. This completes the proof. \square